

Nurse-Led Programs Focusing on Social Support for People with Type 2 Diabetes Mellitus: A Scoping Review*

* This work was conducted with the support of the “Edital N. 21/2018 - Procad/Amazonia,” funding provided by CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior), an agency linked to the Brazilian Ministry of Education, grant number: n.88881.200531/2018-01

Isabela Mendonça Rodrigues dos Santos

<https://orcid.org/0000-0002-1502-7594>
Universidade Federal do Maranhão, Brazil
isabela.mrs@discente.ufma

Isaura Letícia Palmeira Tavares Rolim

<https://orcid.org/0000-0002-8453-2543>
Universidade Federal do Maranhão, Brazil
leticia.isaura@ufma.br

Aurean D’Eça Júnior

<https://orcid.org/0000-0002-7675-412X>
Universidade Federal do Maranhão, Brazil
aurean.junior@ufma.br

✉ **Maria Almira Bulcão Loureiro**

<https://orcid.org/0000-0003-3234-2833>
Universidade Federal do Maranhão, Brazil
almira.maria@discente.ufma.br

Diogo Matheus Barros da Silva

<https://orcid.org/0000-0002-7966-8075>
Universidade Federal do Maranhão, Brazil
silva.diogo@discente.ufma.br

Geysa Santos Góis Lopes

<https://orcid.org/0000-0002-6801-1940>
Rede Sarah Hospitalar, Brazil
geysagois@hotmail.com

Received: 20/06/2023

Sent to peers: 15/08/2023

Approved by peers: 20/10/2023

Accepted: 03/11/2023

DOI: 10.5294/aqui.2024.24.1.2

Para citar este artículo / To reference this article / Para citar este artigo

Santos IMR, Rolim ILTP, D’Eça A, Loureiro MAB, Silva DMB, Lopes GSG. Nurse-led programs focusing on social support for people with type 2 diabetes mellitus: A scoping review. *Aquichan*. 2024;24(1):e2412. DOI: <https://doi.org/10.5294/aqui.2024.24.1.2>

Theme: Care processes and practices

Contribution to the field: This study can contribute to increased adherence to self-care practices and management of this chronic condition. Furthermore, it can promote health education and awareness, adequate glycemic management, promotion of a healthy diet, emotional support, and continuous monitoring by these individuals. Nurse-led programs can result in better adherence and, consequently, long-term prevention.

Abstract

Introduction: Nurse-led programs have a determinant role in the engagement of individuals in the self-management of type 2 diabetes mellitus (T2DM), as they contribute to improving quality of life. Self-management assisted by social support is vital, especially as it describes the importance of nurses in promoting health for people with T2DM, as this intervention can help reduce the consequences and progression of the disease. **Objective:** To map programs focused on nurse-led social support for people with T2DM. **Materials and Methods:** This is a scoping review in line with the methodological guidelines of the Joanna Briggs Institute. The search descriptors used were “*autocuidado*” (self-care), “*intervenção liderada por enfermeiro*” (nurse-led intervention), and “*diabetes mellitus tipo 2*” (type 2 diabetes mellitus), with searches in the Web of Science, Medline, CINAHL, Capes Theses and Dissertations Catalog, Scopus, JBI Evidence Synthesis, PsycINFO, and the Portuguese Open Access Scientific Repositories databases conducted between February and March 2022. The eligibility criteria were studies published within the last 10 years, in English, Spanish, and Portuguese, covering the individual’s experience in programs focused on self-management through nurse-led social support. **Results:** A total of 12 studies were included, all of which were published in the United States, England, and Iran, focusing on social support for self-management in people with T2DM, and centered on care facilities for people with chronic diseases. It was found that most nurse-led programs have been implemented in developed countries and that their main characteristic is their emphasis on social cognitive theory. The population was characterized by being mostly women, aged over 22, with a low level of education. **Conclusions:** The study met the proposed objective by demonstrating that, despite the low implementation cost with high social and economic returns, the development of these programs is still scarce. The need for further studies focusing on emotional levels, such as depression and anxiety, is highlighted.

Keywords (Source: DeCS)

Social Support; Nursing Care; Social Programs; Self-Management; diabetes mellitus type 2.

4 Programas centrados en el apoyo social para personas con diabetes mellitus tipo 2 dirigidos por enfermeros: revisión de alcance*

* Este trabajo fue realizado con el apoyo de “Edital n.21/2018 - Procad/Amazonia”. Financiado por medio de CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior), una agencia vinculada al Ministerio de Educación de Brasil, concedido con el número: n.88881.200531/2018-01

Resumen

Introducción: los programas dirigidos por enfermeras desempeñan un papel fundamental a la hora de implicar a las personas en el autocontrol de la diabetes mellitus tipo 2 (DM2), ya que repercuten en la mejora de la calidad de vida. El autocontrol respaldado por el apoyo social es crucial, especialmente porque describe la importancia de los profesionales de enfermería en la promoción de la salud de las personas con diabetes mellitus tipo 2 (DM2), dado que esta intervención puede ayudar a reducir las consecuencias y el empeoramiento de la enfermedad. **Objetivo:** mapear los programas centrados en el apoyo social para personas con DM2 dirigidos por enfermeros. **Materiales y método:** una revisión de alcance que siguió las directrices metodológicas del Instituto Joanna Briggs. Los descriptores de búsqueda utilizados fueron “autocuidado”, “intervención dirigida por enfermeros”, “diabetes mellitus tipo 2”, con búsqueda en la Web of Science, Medline, CINAHL, Catálogo de Tesis y Disertaciones de la Capes [Coordinación de Perfeccionamiento de Personal de Nivel Superior], Scopus, JBI Evidence Synthesis, PsycINFO y Repositorios Científicos de Acceso Abierto de Portugal, entre febrero y marzo de 2022. Los criterios de elegibilidad fueron estudios publicados en los últimos 10 años, en inglés, español y portugués, que cubrieran la experiencia individual de programas dirigidos por profesionales de enfermería centrados en la autogestión a través del apoyo social. **Resultados:** se incluyeron 12 estudios, publicados en Estados Unidos, Inglaterra e Irán, centrados en el apoyo social para la autogestión de personas con DM2 y en entornos de atención a personas con enfermedades crónicas. Se observó que la mayoría de los programas dirigidos por enfermeros se aplican en países desarrollados y tienen como característica principal el énfasis en la teoría cognitiva social. El público se caracterizó por ser mayoritariamente femenino, mayor de 22 años y con bajo nivel de estudios. **Conclusiones:** el estudio respondió al objetivo propuesto demostrando que, a pesar del bajo costo de implementación con alto retorno social y económico, el desarrollo de estos programas aún es considerado escaso. Se reitera la necesidad de nuevos estudios que enfatizen niveles emocionales como la depresión y la ansiedad.

Palabras clave (Fuente: DeCS)

Apoyo social; atención de enfermería; programas sociales; automanejo; diabetes mellitus tipo 2.

Programas com foco no suporte social para pessoas com diabetes mellitus tipo 2 liderados por enfermeiros: *scoping review**

* Este trabalho foi realizado com apoio do Edital n.21/2018 - Procad/Amazônia. Financiamento via CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior), uma agência vinculada ao Ministério da Educação do Brasil. Financiamento projeto n.88881.200531/2018-01

Resumo

Introdução: os programas liderados por enfermeiros desempenham um papel fundamental no engajamento do indivíduo na autogestão do diabetes *mellitus* tipo 2 (DM2), pois impactam na melhoria da qualidade de vida. A autogestão apoiada pelo suporte social torna-se crucial, sobretudo por descrever a importância do enfermeiro na promoção da saúde da pessoa com DM2, tendo em vista que essa intervenção pode contribuir para reduzir as consequências e os agravamentos da doença. **Objetivo:** mapear os programas com foco no suporte social para pessoas com DM2 liderados por enfermeiros. **Materiais e método:** revisão de escopo que seguiu as diretrizes metodológicas do Joanna Briggs Institute. Utilizou-se como descritores de busca “autocuidado”, “intervenção liderada por enfermeiro”, “diabetes *mellitus* tipo 2”, com busca nos bancos de dados Web of Science, Medline, CINAHL, Catálogo de Teses e Dissertações da Capes, Scopus, JBI Evidence Synthesis, PsycINFO e Repositórios Científicos de Acesso Aberto de Portugal, entre fevereiro e março de 2022. Os critérios de elegibilidade foram estudos publicados nos últimos 10 anos, em inglês, espanhol e português, abrangendo a experiência do indivíduo em programas com foco na autogestão por meio do suporte social liderados por enfermeiros. **Resultados:** 12 estudos incluídos, publicados nos Estados Unidos, na Inglaterra e no Irã, com foco no suporte social de apoio à autogestão de pessoas com DM2 e com concentração em locais de atendimento às pessoas com doenças crônicas. Observou-se que a maioria dos programas liderados por enfermeiros é implementada em países desenvolvidos e apresenta como característica principal ênfase na teoria social cognitiva. O público se caracterizou por ser maioria mulheres, faixa etária de adultos maiores de 22 anos, com baixa escolaridade. **Conclusões:** o estudo respondeu ao objetivo proposto ao demonstrar que, apesar do baixo custo para a implementação com alta devolutiva social e econômica, ainda é considerado escasso o desenvolvimento desses programas. Reitera-se a necessidade de estudos posteriores com ênfase nos níveis emocionais, como depressão e ansiedade.

Palavras-chave (Fonte: DeCS)

Apoio social; cuidados de enfermagem; programas sociais; autogestão; diabetes *mellitus* tipo 2.

Introduction

Type 2 diabetes mellitus (T2DM) represents a public health problem due to its high prevalence, mortality, and morbidity rates, with a resulting increase in costs for the healthcare system. All these negative health outcomes, both at the individual and collective levels, prompt the development of innovative strategies to promote effective self-management (1).

Thus, certain decisions in therapeutic management have an impact on the quality of life of people with T2DM since many of the complications progress to disability. Furthermore, the treatment of complications leads to higher costs, both for the individual and the healthcare system (2). For this reason, the guidelines of scientific organizations dedicated to the study of T2DM recommend that people should participate in education programs for diabetes mellitus self-care (EPDMS). These programs train individuals so that they become knowledgeable enough to make decisions and, simultaneously, induce self-management (3). EPDMS thus benefit and improve this population's health conditions (4).

Self-management is defined as the individual's ability, alongside their families, community, and healthcare professionals, to manage the symptoms, treatment, and psychosocial, cultural, and spiritual changes resulting from chronic conditions. For this purpose, the processes related to self-management include goal setting and self-monitoring, reflective thinking, decision-making, planning and engaging in specific health behaviors, self-assessment, and management of physical, emotional, and cognitive responses regarding health behavior changes (5).

One of the ways to support self-management is by adopting the perspective of social support. Social support is a care network provided by people who are close to an individual or professionals, regardless of where they receive treatment. This form of support provides knowledge and affection, which enables learning, security, and improved self-esteem. These feelings contribute to the continuity of activities regularly, leading to the individual refraining from quitting treatment or withdrawing from society (6).

Those who receive and, especially, perceive the social support provided by their support network can face these adversities with more commitment since they know who to resort to and from whom to receive the necessary help. This helps them to self-manage a pathology's treatment, emphasizing that this is a delicate period, with individual adaptations and resignations (7). From this perspective, interventions based on social support are capable of modifying health conditions, because they reduce the individual's mental and physical stress, as well as reducing behaviors that are harmful to their health (8).

According to the literature, nursing stands out as the profession best suited to provide self-management support, since its impor-

tance lies in the fact that it strengthens the bond for treatment, with knowledge and planning of routines to manage a chronic condition (9). In this sense, specific programs led by nurses that offer social support to teach self-management skills stand out (9). Thus, it is necessary to learn more about nurse-led programs that aim to promote self-management through social support in the field of T2DM, since there is currently a lack of information available on their characteristics. Furthermore, understanding these programs' current state of the art would be useful for informing future research.

Additionally, nursing interventions focused on social support to help the self-management of patients with T2DM contribute to the health-disease process in a way that enables them to manage their health condition. Consequently, it is necessary to carry out this scoping review to synthesize the current scientific evidence on the subject.

The main objective of this review is to map the nature and extent of published scientific production and grey literature related to nurse-led programs aimed at supporting self-management of care of people with T2DM through social support. It has the following specific objectives: To list the programs focusing on social support for people with T2DM led by nurses; to describe the contents of these programs; and to identify the results related to them.

Materials and Methods

This is a scoping review, whose purpose is to map, explore, and describe the literature in the research field and the conceptual limits of the theme addressed. It was done by observing the methodological guidelines of the Joanna Briggs Institute (JBI). Thus, the following steps were followed, according to the Preferred Reporting Items for Systematic Review and Meta-Analyses extension for Scoping Reviews (Prisma-ScR) version 2020: defining and aligning the objectives and questions; developing and aligning the inclusion criteria with the objective and guiding question; describing the planned approach for searching for evidence, selecting and extracting data, and presenting the evidence; searching for the evidence; selecting the evidence; extracting the evidence; evaluating the evidence; presenting the results; summarizing the evidence about the review's purpose, establishing conclusions, and noting any implications arising from the findings (10).

Registration Protocol

The protocol for the preparation of this review is registered on the Open Science Framework (OSF) platform and can be accessed via the following link: <https://osf.io/4wvt6>.

Identification of the Guiding Question

The guiding question formulated was the following: What nurse-led programs exist to support self-management in people with T2DM through social support, and what are their characteristics?

Furthermore, the PCC (population, concept, and context) mnemonic strategy was used to integrate the studies in this review. Therefore, the population consists of adults with T2DM; the concept of nurse-led social support programs; and the context is primary healthcare or the community.

Eligibility Criteria

The eligibility criteria adopted were: studies published with the proposed theme in English, Spanish, and Portuguese; complete articles, theses, and dissertations available online; any study, regardless of methodological design (qualitative, quantitative, or mixed), that explores the participant's experience in programs aimed at supporting the self-management of DM through social support and led by nurses; studies with participants aged 18 or over, diagnosed with T2DM, conducted in the primary health care setting, and with program interventions led by nurses.

Search Strategy

The search for studies was performed in the following databases: Web of Science, Medline (PubMed), Cinahl (EBSCO), Capes Catalog of Theses and Dissertations, Scopus, JBI Evidence Synthesis, PsycINFO (APA), and Open Access Scientific Repositories in Portugal between February 23 and March 17, 2022. Afterward, the reference lists of the included studies were manually searched to identify any relevant studies. The search strategies used for each database are shown in Table 1 and were devised with the assistance of two librarians with expertise in the health field and systematized reviews.

Table 1. Search Strategies Used in the Explored Databases. Brazil, 2022

Database	Search Strategy	Search Date	Retrieved Articles
Cinahl (EBSCO)	((MM "Self Care") OR (MM "Self-Management")) AND (MM "Support, Social/UT") AND (MM "Community Programs") AND (MM "Nurses") AND (MM "Diabetes Mellitus, Type 2").	17/03/2022	11
Open Access Scientific Repositories of Portugal (RCAAP)	"Social support" AND diabetes	17/03/2022	181

Database	Search Strategy	Search Date	Retrieved Articles
Web of Science	("Self Care" OR "self management") AND ("Social Support" OR "Social Care" OR "Online Social Support" OR "Perceived Social Support" OR "Perceived Social Supports") AND ("nurse-led education" OR "nurse-delivered intervention" OR "nurse-led intervention" OR "program development" OR "nurse-led program") AND ("diabetes" OR "diabetes mellitus" OR "diabetes mellitus, Type 2" OR "Diabetes Mellitus, Type II" OR "Type 2 Diabetes Mellitus" OR "Type 2 Diabetes" OR "Diabetes, Type 2")	23/02/2022	3
Scopus	("Self Care" OR "self management") AND ("Social Support" OR "Social Care" OR "Online Social Support" OR "Perceived Social Support" OR "Perceived Social Supports") AND ("nurse-led education" OR "nurse-delivered intervention" OR "nurse-led intervention" OR "program development" OR "nurse-led program") AND ("diabetes" OR "diabetes mellitus" OR "diabetes mellitus, Type 2" OR "Diabetes Mellitus, Type II" OR "Type 2 Diabetes Mellitus" OR "Type 2 Diabetes" OR "Diabetes, Type 2").	23/02/2022	24
PsycINFO	("Self Care" OR "self management") AND ("Social Support" OR "Social Care" OR "Online Social Support" OR "Perceived Social Support" OR "Perceived Social Supports") AND ("nurse-led education" OR "nurse-delivered intervention" OR "nurse-led intervention" OR "program development" OR "nurse-led program") AND ("diabetes" OR "diabetes mellitus" OR "diabetes mellitus, Type 2" OR "Diabetes Mellitus, Type II" OR "Type 2 Diabetes Mellitus" OR "Type 2 Diabetes" OR "Diabetes, Type 2").	23/02/2022	14
JB1 Evidence Synthesis	Self Care AND Social Support AND nurse led AND diabetes Type 2.	09/03/2022	62
Capes Catalog of Theses and Dissertations	"Suporte social" AND "Diabetes"	17/03/2022	1429
PubMed	("Self Care" [MeSH Terms] OR "Self Care" [Text Word] OR "self management" [Text Word]) AND ("Social Support" [MeSH Terms] OR "Social Care" [Text Word] OR "Online Social Support" [Text Word] OR "Perceived Social Support" [Text Word] OR "Perceived Social Supports" [Text Word]) AND ("nurse-led education" [Text Word] OR "nurse-delivered intervention" [All Fields] OR "nurse-led intervention" OR "program development" [All Fields] OR "nurse-led program" [All Fields]) AND ("diabetes" OR "diabetes mellitus" OR "diabetes mellitus, Type 2" OR "Diabetes Mellitus, Type II" OR "Type 2 Diabetes Mellitus" OR "Type 2 Diabetes" OR "Diabetes, Type 2")	09/03/2022	19

Source: Search data (2022).

Study Selection

After searching the databases, all the studies were exported to a reference manager (EndNote). Duplicate articles were removed both manually and using an automation tool. It should be noted that the descriptors were adapted for each database, by using synonyms to find the greatest number of studies.

Next, the titles and abstracts were screened based on the inclusion criteria with the help of the Rayyan Systematic Reviews web/mobile app, so that two authors could independently and blindly screen the title/abstract and full text. In the event of any doubts, a third author would read the manuscript and decide whether or not to include it. Publications that met the eligibility criteria were retrieved in full.

Data Mapping

A form was designed by the author to extract the data. A pretest was then carried out to assess the need for adjustments. The data extracted included author, title, publication year, study country, journal name, document type, language, study objective, study setting, methods, population/sample size, inclusion criteria, exclusion criteria, collection instrument/technique, program results/impacts, conclusions, participants' social characteristics, participants' clinical characteristics, program/intervention duration, core theoretical model, resources used, program implementation, and user satisfaction assessment.

Compiling, Summarizing, and Reporting the Results

A descriptive analysis of the information related to nurse-led programs and their characteristics was carried out. The evidence was synthesized based on the research questions. A table was prepared with data allocated to the studies included in the review and another table with existing programs. Additionally, a figure was prepared with the basic theoretical model of the interventions. The search steps are represented in the flow diagram of the Preferred Reporting Items for Systematic Reviews and Meta-analyses (Prisma-ScR), which is an extension of the scoping review.

Results

Literature Search

Initially, 1,743 publications were identified, as shown in Figure 1. After removing 10 duplicate studies, the titles and abstracts were then screened, resulting in a sample containing 12 documents for full reading. Subsequently, seven publications were excluded for failing to meet the eligibility criteria, which left five publications eligible

for analysis. During the review of the references in the included studies, it was found that it was necessary to incorporate seven additional documents. As a result, this review consisted of 12 studies after the inclusion of the relevant documents.

Most of the selected studies were frequently carried out more than five years ago ($n = 8$) in developed countries such as the United States, England, and Iran ($n = 6$), as shown in Table 2, which presents the main characteristics of the studies included in this review.

Regarding the study designs, there was a prevalence of randomized clinical trials ($n = 8$). The others were systematic reviews plus meta-analysis ($n = 2$), a systematic review ($n = 1$), and a longitudinal observational study ($n = 1$). As for document type, eight were articles, three were theses, and one was a dissertation.

Study Characteristics


Nurse-led programs were implemented in the following locations: general care clinics ($n = 6$), specialized care services for people with chronic illnesses ($n = 3$), and university hospitals ($n = 3$), as shown in Table 3. It is noted that the programs are designed to improve glycemic control and, simultaneously, achieve significant results in terms of lipid profile, blood pressure, regular physical activity, adequate nutrition, emotional stability, social interaction and, consequently, self-management.

The participants' characteristics indicate a mean age of 54.2 ± 11.8 years (between 22 and 69 years), with a prevalence of women ($n = 7$), with a low level of education, having studied for a total of nine years ($n = 8$).

The studies found on emotional support covered programs consisting of individual, face-to-face counseling, followed by a collective approach and dynamics with family/friends. To consolidate learning, phone calls were made (Table 4). Therefore, the programs converged on an educational basis—with instructions in face-to-face appointments and via phone calls—with a strong influence of affective aspects, since the person with diabetes would need to be the protagonist of their decisions, but there was also family interaction. On average, the interventions lasted 12 months, with a maximum of 36 months and a minimum of 1 month.

Regarding the basic theoretical model used by the programs, most followed the social cognitive theory ($n = 3$), conversation maps ($n = 2$), the common sense self-regulation theory ($n = 2$), the self-efficacy theory ($n = 1$), and motivational interviewing ($n = 1$). It is worth noting that the studies that were not mentioned correspond to the fact that they did not present their respective basic models and/or clarified the theoretical framework.

Figure 1. Study identification and inclusion process. Prisma-ScR flowchart



Source: Own elaboration.

Table 2. Studies included in the scoping review by author, type of study, title, country, and year of publication. Brazil, 2022

Author	Type of Study	Title	Country	Year of Publication
Klein HA, Jackson SM, Street K, Whitacre JC, Klein G	Article: systematic review + meta-analysis	Diabetes self-management education: Miles to go	United States	2012
Pereira DA, Costa NMSC, Sousa ALL, Jardim PCBV, Zanini CRO	Article: randomized controlled clinical trial	Efeito de intervenção educativa sobre o conhecimento da doença em pacientes com diabetes mellitus	Brazil	2012
Parker D, Clifton K, Shams R, Young J	Article: systematic review	The effectiveness of nurse-led care in general practice on clinical outcomes in adults with type 2 diabetes	Australia	2012
Mahdizadeh M, Peyman N, Thagipour A, Esmaily H	Article: randomized controlled clinical trial	Effect of health education program on promoting physical activity among diabetic women in Mashhad, Iran: Applying social cognitive theory	Iran	2013
Bertolin DC	Thesis: randomized controlled clinical trial	Estresse, modos de enfrentamento e aceitação da doença de pessoas com diabetes mellitus tipo 2 participantes de um programa educativo como envolvimento de um familiar: ensaio clínico randomizado	Brazil	2013
Tang PC, Overhag JM, Chan AS, Brown NL, Aghighi B, Entwistle MP et al.	Article: clinical trial controlado e randomizado	Online disease management of diabetes: Engaging and motivating patients online with enhanced resources-diabetes (EMPOWER-D), a randomized controlled trial	The United States	2013
Villas Boas LCG, Pace AE	Thesis: randomized controlled clinical trial	Contribuição do apoio social familiar nos resultados das intervenções educativas junto às pessoas com diabetes mellitus tipo 2: ensaio clínico controlado randomizado	Brazil	2014
Gomides DS	Dissertation: longitudinal observational study	Resultados de intervenções para o cuidado do diabetes mellitus com foco no apoio social: estudo longitudinal	Brazil	2016
Murray E, Sweeting M, Dack C, Pal K, Modrow K, Hudda M et al.	Article: randomized controlled clinical trial	Web-based self-management support for people with type 2 diabetes (HeLP-Diabetes): Randomised controlled trial in English primary care	England	2017
Golnaz A, Kim LS, Shariff GS, Said S, Sanaz A, Mosayeb M et al.	Article: double-blind randomized clinical trial	Effect of a nurse-led diabetes self-management education program on glycosylated hemoglobin among adults with type 2 diabetes	Iran	2018
Hadjiconstantinou M	Thesis: systematic review and meta-analysis + focus groups	The development of an online emotional support programme, to improve diabetes related distress and well-being in type 2 diabetes	England	2018
Van Puffelen AL, Rijken M, Heijmans MJWM, Nijpels G, Schellevis	Article: randomized clinical trial	Effectiveness of a self-management support program for type 2 diabetes patients in the first years of illness: Results from a randomized controlled trial	The Netherlands	2019

Source: Research data (2022).

Table 3. Nurse-led programs focusing on social support for people with type 2 diabetes by author, intervention site, results achieved, and population assisted.

Author	Intervention Site	Intervention Results	Population Assisted
Klein HA, Jackson SM, Street K, Whitacre JC, Klein G	Randomly assigned	A significant reduction in blood glyated hemoglobin levels.	Age (mean): ND Sex: ND Education: ND
Pereira DA, Costa NMSC, Sousa ALL, Jardim PCBV, Zanini CRO	Referral service for hypertension treatment	Increased knowledge of diabetes mellitus.	Age (mean): 65.8 years Sex: Female Education: 8 years of education
Parker D, Clifton K, Shams R, Young J	General/family practice environment	Improvement in clinical parameters: Changes in systolic and diastolic blood pressure (BP), body mass index (BMI), cholesterol levels, HbA _{1c} (glycated hemoglobin), and fasting glycemia.	Age: >18y Sex: ND Education: ND
Mahdizadeh M, Peyman N, Thagipour A, Esmaily H	Healthcare centers in Mashhad City, eastern Iran	In the intervention group (IG), light physical activity and sedentary behavior decreased from 56.1 % (23 individuals) to 14.6 % (6 individuals) in the next stage. There was a significant improvement over time in terms of physical activity (increase in the mean number of minutes practiced per week). Adherence to physical activity was higher in the groups due to the social support received, mainly from family, friends, and exercise partners.	Age (mean): 48.37 years Sex: Female Education: ND
Bertolin DC	Diabetes outpatient clinic at a tertiary level healthcare university hospital unit	The most commonly mentioned coping methods are cognitive restructuring, information-seeking, medium acceptance of the disease, low perception of stress, and some level of stress (in the resistance stage of the disease).	Age (mean): 60.4 years Sex: Female Education: A minimum of 0 and maximum of 17 years of education
Tang PC, Overhag JM, Chan AS, Brown NL, Aghighi B, Entwistle MP et al.	Palo Alto Medical Foundation, a non-profit health organization	The IG had significantly better diabetes management than the control group (CG) in the first 6 months. There was no significant difference after 12 months. The IG had significantly better management of LDL cholesterol at 12 months. There was no statistically significant difference in BP management, weight, and Framingham risk at 12 months.	Idade: > 18 years Sex: Male Education: >12 years of education

Author	Intervention Site	Intervention Results	Population Assisted
Villas Boas LCG, Pace AE	Hospital das Clínicas da Faculdade de Medicina de Ribeirão Preto da Universidade de São Paulo (HCFMRP-USP)	<p>There were no statistically significant differences after carrying out the interventions for perceived self-efficacy, knowledge about diabetes, medication adherence, and perceived social support.</p> <p>For the self-care variable, the CG performed better than the IG in one behavior related to eating and in two related to physical activity.</p> <p>Regarding the clinical variables, the CG had a lower body mass index and lower abdominal circumference in women.</p> <p>Metabolic management: there were no statistically significant differences, but there was a reduction in glycated hemoglobin in favor of the intervention group.</p>	<p>Age (mean): 60.43 years Sex: Female Education: < 9 years</p>
Gomides DS	Outpatient clinic for diabetes treatment at a tertiary care hospital	<p>There were no differences between the groups for the variables BP, waist circumference, BMI, glycated hemoglobin, fasting plasma glucose, self-care activities, knowledge about the disease, and self-efficacy.</p> <p>The timeframe was defined as baseline, the end of the interventions, and 24 months after the interventions. When comparing the times, there was a reduction in BP and glycated hemoglobin and an increase in BMI and abdominal circumference in both groups.</p> <p>Self-care knowledge and activities increased immediately after the interventions had been carried out and decreased after two years in both groups.</p> <p>Self-efficacy was maintained in the IG and reduced in the CG after two years and at the end of the interventions.</p> <p>Interventions including social support had no difference between the groups and over time, through descriptive analysis, there was an improvement in self-efficacy, which was sustained after two years in the IG.</p> <p>Group educational interventions were effective in improving glycemic control and BP two years after the interventions had been carried out.</p> <p>For the knowledge variables about the disease and self-care, the results suggest the need for educational reinforcement over time.</p>	<p>Age: from 45 to 84 years Sex: Female Education: From 0 to 17 years of education</p>

Author	Intervention Site	Intervention Results	Population Assisted
Murray E, Sweeting M, Dack C, Pal K, Modrow K, Hudda M et al.	21 general practices in England combining urban, suburban, and rural practices	<p>The IG participants had lower glycosylated hemoglobin (HbA_{1c}) than those in the CG.</p> <p>There were no significant overall differences between the groups in the diabetes-related stress score, measured by the problem areas in diabetes (PAID) scale. However, for participants who had been diagnosed with diabetes more recently, it indicated a beneficial impact of the intervention on this group.</p>	<p>Age: 65 years</p> <p>Sex: Male</p> <p>Education: ND</p>
Golnaz A, Kim LS, Shariff GS, Said S, Sanaz A, Mosayeb M et al.	Urban primary and secondary endocrine outpatient clinic located in a university hospital in the city of Ilam, Iran	The IG patients had a significant improvement in glycosylated hemoglobin, BP, body weight, efficacy expectancy, outcome expectancy, and diabetes self-management behaviors.	<p>Age (mean): 54.2 ± 11.8 years (ranging from 22 to 69 years)</p> <p>Sex: Female</p> <p>Education: 9 years of education</p>
Hadjiconstantinou M	Leicester Diabetes Centre	<p>Online support led by healthcare professionals was associated with positive well-being outcomes compared to those with non-professional support.</p> <p>Individually, several studies found significant improvements in well-being measurements. This was not supported by the meta-analysis for the outcome measurements of depression and distress.</p>	<p>Age (mean): 53.4 years</p> <p>Sex: Female</p> <p>Education: ND</p>
Van Puffelen AL, Rijken M, Heijmans MJWM, Nijpels G, Schellevis	General practice clinics in the Netherlands.	<p>Immediately after the program, the IG presented a significantly greater increase in physical activity and fruit and vegetable intake. Six months after the intervention, these effects had disappeared.</p> <p>Regarding diabetes-related suffering, no immediate or six-month effects were found.</p> <p>The baseline levels of self-care seemed to be slightly higher in the IG, but not significantly higher.</p> <p>Six months after the intervention, the differences in empowerment were still present, even though healthy behaviors had decreased.</p> <p>Considering distress, there were no significant changes, as levels of distress were low even before the intervention had been carried out.</p>	<p>Age (mean): 63.5 years</p> <p>Sex: Male</p> <p>Education: 12 years of education</p>

Source: Research data (2022).

Key: ND — not defined.

Table 4. Programs by Author, Type of Intervention, and Duration

Author	Type of Intervention	Duration
Klein HA, Jackson SM, Street K, Whitacre JC, Klein G	<p>1. Rules and procedures: Rules concerning diet and exercise. Rules for procedures, such as using a diary to record daily glucose levels.</p> <p>2. Affective: Encouragement and motivation to build trust.</p> <p>3. Social and situational: Approach focused on managing social and situational factors. Strategies that encompass planning meals on vacation and in restaurants.</p> <p>4. Complex cognition: Use of mental models or other cognitive strategies to support the understanding of diabetes, which are used to moderate blood glucose levels.</p>	<p>Three categories: 13 weeks or less, 14-26 weeks, 27 weeks or more.</p>
Pereira DA, Costa NMSC, Sousa ALL, Jardim PCBV, Zanini CRO	<p>The IG participated in 12 educational meetings over six months, held every two weeks and lasting an average of two hours. The first two educational meetings were held to identify the learning needs and previous knowledge about the disease among the research participants.</p> <p>A pre-established script was used for the discussion, which aimed to identify the patients' knowledge of diabetes, the difficulties caused by the disease, and their perception of what could help with treatment.</p> <p>Organization of educational activities: a teaching plan was devised. Each plan used distinctive resources, such as pictures, phrases written by the patients themselves, posters, songs, walks, questions, and verbal expressions of meaningful experiences.</p>	6 months
Parker D, Clifton K, Shams R, Young J	Care is guided by nurses who are recognized as having a leadership role in the care of patients with T2DM compared to care provided by general practitioners.	Not defined.
Mahdizadeh M, Peyman N, Thagipour A, Esmaily H	<p>The training program has been introduced. Introductory session: The rules have been announced. First session: Discussion regarding a healthy lifestyle. Second session: Discussion about problems and the search for alternatives to overcome barriers. Third session: Assigning objectives and their importance in achieving favorable results. Fourth session: Dialogue about having a regular program and planning physical activity. Fifth, sixth, and seventh sessions: Walking program and practical training course in physical activity.</p>	7 sessions (4 theoretical and 3 practical) in 1 month (each session ≈ 60 min)
Bertolin DC	<p>Intervention with people with diabetes: performed in groups, using conversation maps. It is a tool that aims to empower people to make decisions. It provides discussions and dynamics in groups of up to 10 people. Maps were used, focusing on how the body and diabetes work (map 1), healthy eating and physical activity (map 2), monitoring blood glucose (map 3), and achieving insulin-related goals (map 4). A total of four meetings were held.</p> <p>Intervention with family members of people with diabetes: Through phone calls. Weekly calls were made, using the motivational interviewing technique. In this technique, the professional establishes an empathetic and collaborative relationship with the person. It follows pre-established protocols and uses the social cognitive theory. The topic of the conversation maps from the group meetings is addressed. A total of 17 calls were made.</p>	23 months

Author	Type of Intervention	Duration
Tang PC, Overhag JM, Chan AS, Brown NL, Aghighi B, Entwistle MP et al.	1st) Home glucose meter readings with graphical feedback; 2nd) Comprehensive report of the patient's specific diabetes status; 3rd) Nutritional and workout records; 4th) Insulin record; 5th) Online messaging with the patient's healthcare team; 6th) Nursing care manager and nutritionist providing advice and medication management, and 7th) Short personalized texts and educational videos provided electronically by the healthcare team.	12 months
Villas Boas LCG, Pace AE	<p>IG: Educational intervention with the participation of the family caregiver through phone calls. Returns occur every three months. A second assessment was done at 12 months. CG: No family member participation.</p> <p>Group educational interventions for people with T2DM: A maximum of eight participants. Approach carried out based on diabetes mellitus conversation maps. This tool aims to empower the individual to self-manage. The topics covered followed those available in Portuguese (how the body and diabetes work; healthy eating and physical activity; medication treatment; blood glucose monitoring; and achieving insulin-related goals).</p> <p>Educational interventions for family caregivers of people with T2DM: Phone calls following motivational interviews with a technique designed to generate collaborative empathy. Topics previously established by the diabetes conversation maps were discussed. A total of 17 calls lasting an average of eight minutes each have been made.</p>	Approximately 12 months
Gomides DS	Educational interventions that provide information related to the disease and the self-care activities required for treatment, using visual and interactive tools based on the social cognitive theory, whose key concept is personal belief in one's ability to perform certain tasks (self-efficacy).	The study times were baseline (T ₀), end of interventions (T ₁₂), and 24 months after completion (T ₃₆).
Murray E, Sweeting M, Dack C, Pal K, Modrow K, Hudda M et	<p>It is an <i>application</i> developed specifically for people with T2DM.</p> <p>The contents present in this application include information about diabetes, how diabetes is treated, possible complications from diabetes, possible impacts of diabetes on relationships at home and work, dealing with unusual situations such as parties, holidays, travel, or shift work, and what lifestyle changes will improve health, skills, and behavior change.</p> <p>Special modules cover behavior change, healthy eating, weight loss, being more physically active, quitting smoking, moderating alcohol consumption, managing medication, glycemic management, and BP management.</p> <p>In addition, content on emotional well-being and sharing experiences between users is available.</p> <p>Possibility of receiving monthly notifications with relevant news about diabetes.</p>	12 months

Author	Type of Intervention	Duration
Golnaz A, Kim LS, Shariff GS, Said S, Sanaz A, Mosayeb M et al.	<p>The participants in the intervention group received a booklet at the beginning of the program, whose content was used throughout the intervention to guide learning, development, and discussion or it was used to create resources for self-guided learning.</p> <p>Within the first four weeks of the intervention, participants were invited to watch four weekly 10-minute films on the prevention of short- and long-term complications (session 1), physical activity, daily foot care (session 2), healthy eating (session 3), and healthy living with diabetes (session 4).</p> <p>Four group discussion sessions were held at weekly intervals, but the schedule was flexible, and attendance was limited to 10 participants for each session. These sessions were held weekly during the first four weeks of the intervention. Each session lasted 120 minutes. All group members were encouraged to attend each session. Group members who missed a session received a follow-up from the moderator before the next session. At the last meeting of the group session, all group members were handed a list of home phone numbers and encouraged to ask for help if necessary.</p> <p>Two months following the end of the group discussion sessions, the intervention participants received a phone call once a week. Each follow-up call lasted approximately 15 to 20 minutes.</p> <p>At the end of the intervention, the CG participants were handed an educational booklet and films.</p>	3 months
Hadjiconstantinou M	<p>Techniques for behavior change: To provide information; tracking/self-monitoring; to provide motivation; to provide feedback; goal setting; problem-solving; action planning; social support; emotional regulation training, and immediate review of behavioral goals.</p>	A minimum of 6 months and a maximum of 18 months.
Van Puffelen AL, Rijken M, Heijmans MJWM, Nijpels G, Schellevis	<p>It consists of three monthly two-hour interactive sessions and a follow-up session three months after the last session.</p> <p>The sessions were led by two nurses specializing in diabetes or nurses who received four hours of training and a detailed manual describing the content of the program and its underlying theories to be used during the sessions.</p> <p>The participating patients and their partners received a <i>booklet</i> containing basic information on diabetes, (homework) assignments, and theoretical and practical information on the topics discussed throughout the course (both books are available in Dutch from the author).</p> <p>The CG participants were invited to a single two-hour educational lecture, in which they received information about living with diabetes from a medical perspective. A professor of general practice and diabetes care provided information about the course of diabetes (including treatment options and possible complications) and the latest developments in diabetes research, according to the classic didactic method.</p>	8 months

Source: Research data (2022).

Evidence Summary

The present study found studies conducted in England, the United States, Iran, Australia, and the Netherlands on programs focused on social support for the self-management of people with T2DM, which were led by nurses and centered on care facilities for people with chronic diseases (9, 11-15).

It is worth noting that in these countries, care for people with diabetes is performed in a unique way and with a holistic approach (16). In developing countries such as Brazil, this approach is still in its infancy, despite efforts to improve care for this population, which has increased significantly over the years (17). Factors such as adequate spending of financial resources on healthcare services, continued training processes and engagement of professionals, as well as technological support and priority on conflict management may be associated with the differences mentioned between these regions (18-20).

In addition, strategies for patients' attendance and commitment to the treatment of chronic diseases are related to family engagement and effective professional intervention, which highlights the importance of social support in achieving these goals. Such support is an essential tool for promoting improvements in health and interpersonal relationships, which leads to improved cognitive performance, and negative emotional reactions management —e.g., anxiety, and offers behavioral analysis (21, 22). The literature highlights the key role played by nursing professionals in primary care in health education activities and self-management through evidence-based practices (23).

In the present study, the findings regarding the theoretical framework that integrates social support into the main strategic interventions were the social cognitive theory (an active pedagogical strategy that determines that individuals influence and are influenced by observational learning [13, 24-26]); conversation maps (a tool subdivided into four stages, delivered to patients via images and metaphors reproduced on a card table with specific topics [27]); the self-efficacy theory (which determines that commitment, depending on personal will and ability, is decisive in preventing or managing the disease [28]); motivational interviewing (evidence-based counseling, in which healthcare professionals help people adopt targeted treatment recommendations [29]); and the common sense self-regulation theory (which provides strategies aimed at finding out how people feel and think about the disease [30]).

These theories can lead to several benefits for patients, such as a reduction in sedentary lifestyles, which has an impact on BMI control and abdominal circumference in women, a better understanding

and acceptance of the disease, as well as stress reduction and increased consumption of fruit and vegetables (13, 24, 25). Thus, knowledge of these theoretical models can enable them to be implemented through low-cost programs and favor care processes with a reduction of risks and problems (31, 32).

A study published in the *Revista de Medicina da Universidade de São Paulo* showed that the perception of social support and family functionality were positive and there was a significant relationship between glycemic levels and information on the adequate management of T2DM in elderly males, which highlights the importance of family support in the therapeutic management of the disease. Nurse-led programs focused on promoting social support and family integration can promote better engagement between patients and their families and, consequently, contribute to improved self-care, healthy eating habits, and physical activity (33-35). In addition, it is worth highlighting the inclusion of people with T2DM in support groups and educational activities, as this allows for joint interaction, and sharing experiences and feelings (36).

Regarding the multidisciplinary approach, nutritional interventions focused on fruit and vegetable intake and physical exercise programs were noted (15, 25, 37). In the field of nursing, the use of online tools was the main form of intervention and satisfactory results were achieved in terms of glycosylated hemoglobin values, fasting glycemia, BP levels, and anthropometric data (9, 14, 24, 31, 37-39, 44). It is worth pointing out that, in the field of primary care, nutritional monitoring associated with an adequate diet has led to the best short-term results with a balanced diet specific to each individual (40).

Regarding physical training programs for people with T2DM, these provided greater benefits by improving insulin sensitivity and glycosylated hemoglobin, which resulted in lower postprandial hyperglycemic peaks. The recommendation to practice physical exercise between 150 and 300 minutes (light to moderate) or 75 to 150 minutes (intense) has an impact on muscle strength and greater endurance capacity in these individuals, which can improve functional capacity and body maintenance (41). In addition, the professional support provided to recently diagnosed patients has benefits, such as reducing stress levels, especially in those who have been diagnosed within less than five years, and who are more resistant to accepting the disease (13, 15, 31, 44).

However, it is worth noting that, although self-management is effective in changing the cognition of people with T2DM, contributing to behavioral change and emotional perception, the maintenance of these behaviors still seems to be insufficient in the long term; therefore, an approach with individuals who have been diagnosed for longer is also essential, as it ensures continuity of care, reduces the possibility of chronic complications such as neuropathy, retinopathy, renal and neurological damage (25, 42).

The present review found that online programs and phone-based approaches under the guidance and leadership of a nurse were more likely to lead to adherence by participants. This can be justified by the fact that technological resources make communication easier, although access to these devices must be considered (9, 12-15, 24, 25, 31, 39, 43, 44).

Thus, due to the complexity of the treatment, effective adherence by patients becomes a challenge not only for them but also for the healthcare team and family members. Consequently, comprehensive support and reframing of these individuals' lifestyle habits must have an impact to promote the practice of effective self-care, as well as conscious decision-making with empowerment through cognitive restructuring (25, 44-47).

As such, the social importance of nurse-led programs for individuals with T2DM is materialized as an important alternative in the follow-up of these individuals in terms of risk reduction, adequate self-care management, and satisfactory family support, which contributes to personal satisfaction and lower demand for healthcare services (9, 48-50).

Limitations and Strengths

Although the methodological evaluation of studies is not a stage covered by the scoping review methodology, this can influence the results. The study was limited to the English, Portuguese, and Spanish languages, which may represent a selection bias. Thus, programs from other languages/cultures are not represented in this study. In addition, the studies did not clarify the follow-up and monitoring of the participants included in them, which hinders identifying and characterizing them in the medium and long term.

The strengths of the review include the rigorous and transparent approach adopted to select relevant studies, conducted in line with the Prisma-ScR guidelines. The strengths also lie in the collaboration of a multidisciplinary team of reviewers and the collaboration of librarians with expertise in systematic and scoping reviews to devise the search string.

Conclusions

Based on the results presented, it was found that nurse-led social support programs for self-management are patient- and family-centered and also use playful and educational auditory or visual strategies and can be based on collective and face-to-face sessions and phone calls. The main results of these programs were improved glycemic control, hemodynamics, and lipid profile. In addition, there has been an improvement in emotional and behavioral parameters, especially in the perception of stress and self-management.

Although there is a tendency for an increase in the number of these programs, there are still gaps that warrant attention; there is a need for future studies focused on emotional levels, such as depression and anxiety, as there was no unanimity on these findings. These psycho-emotional aspects can negatively influence the physiological and metabolic parameters of patients with T2DM. Furthermore, there is a need to conduct studies with individuals who have been diagnosed for longer, as there was a prevalence of individuals who had been diagnosed less than five years ago.

Nurse-led programs focusing on social support seem to be an intervention that can lead to more effective self-management, meeting people's real needs and, consequently, achieving positive health outcomes in the T2DM setting. The relevance of this type of study in following up with participants in the medium and long term is emphasized to highlight the benefits of the proposed interventions.

Conflicts of interest: None declared.

References

- International Diabetes Federation. IDF diabetes atlas: 10th edition. International Diabetes Federation; 2021. p. 905-911.
- Caires JM, Oliveira JC, Simonetti SH, Leal MR, Marreira M. The role of nurses as health educator so patients with Diabetes Mellitus type 2: Integrative review. RSD [Internet]. 2022;11(12):e487111234726. DOI: <https://doi.org/10.33448/rsd-v11i12.34726>
- Joaquim N, Nunes T, Miranda R. Intervenções educacionais baseadas na família para a autogestão da diabetes em adultos: revisão baseada na evidência. Rev Port Med Geral Fam [Internet]. 2022;38(1):61-72. DOI: <https://doi.org/10.32385/rpmgf.v38i1.13064>
- Marques IC. Diabetes Mellitus: principais aspectos e diagnóstico através da dosagem de hemoglobina glicada. (monografia de graduação em farmácia). Ouro Preto: Universidade Federal de Ouro Preto; 2018. Disponível em: <http://www.monografias.ufop.br/handle/35400000/1096>
- Barbara R, Heleen W, Paolo IDE, Ingrid BC, Joyce BS, Michael ASB et al. Characteristics of self-care interventions for patients with a chronic condition: A scoping review. International journal of nursing studies. 2021;116:103713. DOI: <https://doi.org/10.1016/j.ijnurstu.2020.103713>
- Ramkisson S, Pillay BJ, Sibanda W. Social support and coping in adults with type 2 diabetes. African J Primary Health Care Fam Med. 2017;9(1):1-8. DOI: <https://doi.org/10.4102/phcfm.v9i1.1405>
- Nunes TGR, Pontes FAR, Silva LIC. Juventude e apoio social: um olhar sobre as redes sociais de estudantes paraenses. Práxis Educativa. 2020;15:1-21. DOI: <https://doi.org/10.5212/PraxEduc.v15.13534.017>
- Poblete F, Barticevic N, Sapag JC, Tapia P, Bastías G, Quevedo D et al. Social support, self-rated health, treatment adherence and effectiveness in patients with type II diabetes and hypertension. Rev Med Chile [Internet]. 2018;146(10):1135-42. DOI: <https://doi.org/10.4067/S0034-98872018001001135>
- Golnaz A, Kim LS, Shariff GS, Said S, Sanaz A, Mosayeb M et al. Effect of a nurse-led diabetes self-management education program on glycosylated hemoglobin among adults with type 2 diabetes. Journal of diabetes research. 2018;8:4930157. DOI: <https://doi.org/10.1155/2018/4930157>
- Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, Moher D et al. Prisma extension for Scoping Reviews (Prisma-ScR): Checklist and explanation. Annals of internal medicine. 2018;169(7):467-73. DOI: <https://doi.org/10.7326/M18-0850>
- Pereira DA, Costa NMSC, Sousa ALL, Jardim PCBV, Zanini CRO. The effect of educational intervention on the disease knowledge of diabetes mellitus patients. Rev Latino-Americana de Enfermagem. 2012;20:478-85. DOI: <https://doi.org/10.1590/S0104-11692012000300008>
- Bertolin DC. Estresse, modos de enfrentamento e aceitação da doença de pessoas com diabetes mellitus tipo 2 participantes de um programa educativo com o envolvimento do familiar: ensaio clínico randomizado. (tese de doutorado em ciências). São Paulo: Universidade de São Paulo; 2013. DOI: <https://doi.org/10.11606/T.22.2013.tde-22012014-110133>
- Gomides DS. Resultados de intervenções para o cuidado do diabetes mellitus com foco no apoio social: estudo longitudinal. (dissertação de mestrado em ciências). São Paulo: Universidade de São Paulo; 2016.
- Hadjiconstantinou M. The development of an online emotional support programme to improve diabetes-related distress and well-being in type 2 diabetes. (tese de doutorado em Filosofia). Leicester: University of Leicester; 2018.
- Van Smoorenburg AN, Hertroijs DFL, Dekkers T, Melles M. Patients' perspective on self-management: Type 2 diabetes in daily life. BMC Health Serv Res. 2019;19:605. DOI: <https://doi.org/10.1186/s12913-019-4384-7>
- Lavorato GN, Benevides SSN, Parisi MCR. Teaching strategies of clinical management of DM2 in medical undergraduate degree: Integrative review. Teaching Strategies of Clinical Management of DM2 in medical undergraduate degree: Integrative review.

- SciELO Preprints. 2022. DOI: <https://doi.org/10.1590/SciELOPreprints.4534>
17. Mor N. Lessons for developing countries from outlier country health systems. *Frontiers in public health*. 2022;10. DOI: <https://doi.org/10.3389/fpubh.2022.870210>
 18. Catic T, Popovic SP, Asimi ZV, Hlavinkova L. Costs of diabetes mellitus and its complications in Bosnia and Herzegovina. *Mater Sociomed*. 2022;34(2):149-54. DOI: <https://doi.org/10.5455/msm.2022.34.149-154>
 19. Trankle SA, Usherwood T, Penelope A, Mary R, Michael C, Christian MG et al. Key stakeholder experiences of an integrated healthcare pilot in Australia: A thematic analysis. *BMC Health Services Research*. 2020;20:925. DOI: <https://doi.org/10.1186/s12913-020-05794-2>
 20. Lynch MK, Bazier A, Ward S, Rand KL, Hirsh AT. Resilience, social support, and health in emerging adults with and without chronic health conditions. *Emerging Adulthood*. 2023;11(3):557-71. DOI: <https://doi.org/10.1177/21676968221148322>
 21. Miao Jonasson J, Hendryx M, Shadyab AH, Kelley E, Johnson KC, Kroenke CH et al. Social support, social network size, social strain, stressful life events, and coronary heart disease in women with type 2 diabetes: A cohort study based on the Women's Health Initiative. *Diabetes care*. 2020;43(8):1759-66. DOI: <https://doi.org/10.2337/dc19-2065>
 22. Kumah E, Afriyie EK, Abuosi AA, Ankomah SE, Fusheini A, Otchere G. Influence of the model of care on the outcomes of diabetes self-management education program: A scoping review. *J Diabetes Res*. 2021;2969243. DOI: <https://doi.org/10.1155/2021/2969243>
 23. Villas Boas LCG, Pace AE. Contribuição do apoio social familiar nos resultados das intervenções educativas junto às pessoas com diabetes mellitus tipo 2: ensaio clínico controlado randomizado (tese de doutorado em ciências). Ribeirão Preto: Escola de Enfermagem de Ribeirão Preto, Universidade de São Paulo; 2014. DOI: <https://doi.org/10.11606/T.22.2014.tde-07012015-141734>
 24. Van Puffelen AL, Rijken M, Heijmans MJWM, Nijpels G, Schellevis FG. Diacourse study group. Effectiveness of a self-management support program for type 2 diabetes patients in the first years of illness: Results from a randomized controlled trial. *PLoS One*. 2019;14(6):e0218242. DOI: <https://doi.org/10.1371/journal.pone.0218242>
 25. Lima CR, Menezes IHCF, Peixoto MDRG. Educação em saúde: avaliação de intervenção educativa com pacientes diabéticos, baseada na teoria social cognitiva. *Ciência & Educação (Bauru)*. 2018;24(1):141-56. DOI: <https://doi.org/10.1590/1516-731320180010010>
 26. Defeudis G, Khazrai YM, Di Rosa C, Secchi C, Montedoro A, Maurizi AR et al. Conversation Maps™, an effective tool for the management of males and females with type 2 diabetes and mildly impaired glycemic control. *Hormones (Athens, Greece)*. 2018;17(1):113-7. DOI: <https://doi.org/10.1007/s42000-018-0005-9>
 27. Guise S. Mini hábitos: cómo lograr grandes resultados con el mínimo esfuerzo. *Málaga: Editorial Sirio*; 2016.
 28. Tosun A, Zincir H. The effect of a transtheoretical model-based motivational interview on self efficacy, metabolic control, and health behaviour in adults with type 2 diabetes mellitus: A randomized controlled trial. *International Journal of Nursing Practice*. 2019;25(4):e12742. DOI: <https://doi.org/10.1111/ijn.12742>
 29. Xin M, Chan VWY, Kong APS, Lau JTF, Cameron LD, Mak WWS et al. Using the common-sense model to explicate the role of illness representation in self-care behaviours and anxiety symptoms among patients with Type 2 diabetes. *Patient Education and Counseling*. 2023;107:107581. DOI: <https://doi.org/10.1016/j.pec.2022.107581>
 30. Murray E, Sweeting M, Dack C, Pal K, Modrow K, Hudda M et al. Web-based self-management support for people with type 2 diabetes (HeLP-Diabetes): Randomised controlled trial in English primary care. *BMJ open*. 2017;7(9):e016009. DOI: <https://doi.org/10.1136/bmjopen-2017-016009>
 31. Subramanian SC, Porkodi A, Akila P. Effectiveness of nurse-led intervention on self-management, self-efficacy and blood glucose level among patients with Type 2 diabetes mellitus. *Journal of complementary & integrative medicine*. 2020;17(3). DOI: <https://doi.org/10.1515/jcim-2019-0064>
 32. Sousa-Munoz RL, Sá AD. Apoio social, funcionalidade familiar e controle glicêmico de pacientes diabéticos tipo 2. *Rev Med*. 2020;99(5):432-41. DOI: <https://doi.org/10.11606/issn.1679-9836.v99i5p432-441>
 33. Kim EJ; Han KS. Factors related to self-care behaviours among patients with diabetic foot ulcers. *Journal of Clinical Nursing*. 2020;29(9-10):1712-22. DOI: <https://doi.org/10.1111/jocn.15215>
 34. Cardoso SB, Amorim FCM, Silva SMS, Carvalho MCA, Pereira GF, Carvalho CMS et al. Atuação do enfermeiro na prevenção e controle do Diabetes mellitus tipo 2. *Research, Society and Development*. 2022;11(13) e139111334563. DOI: <https://doi.org/10.33448/rsd-v11i13.34563>
 35. Moreira JB, Muro ES, Monteiro LA, Lunes DH, Assis BB de, Chaves E de CL. Efeito do grupo operativo no ensino do autocuidado com os pés de diabéticos: ensaio clínico randomizado. *Rev esc enferm USP [Internet]*. 2020;54:e03624. DOI: <https://doi.org/10.1590/s1980-220x2019005403624>
 36. Klein HA, Jackson SM, Street K, Whitacre JC, Klein G. (2013). Diabetes self-management education: miles to go. *Nursing Research and Practice*. 2013;581012. DOI: <https://doi.org/10.1155/2013/581012>
 37. Parker D, Clifton K, Shams R, Young J. The effectiveness of nurse-led care in general practice on clinical outcomes in adults with type 2 diabetes. *JBI library of systematic reviews*. 2012;10(38):2514-58. DOI: <https://doi.org/10.11124/jbisrir-2012-33>
 38. Tang PC, Overhag JM, Chan AS, Brown NL, Aghighi B, Entwistle MP et al. Online disease management of diabetes: Engaging and motivating patients online with enhanced resources-diabetes (EMPOWER-D): A randomized controlled trial. *Journal of the American Medical Informatics Association*. 2013;20(3):526-34. DOI: <https://doi.org/10.1136/amiainl-2012-001263>
 39. Namazi N, Esmaeili S, Ahmadikhatir S, Razi F, Nasli-Esfahani E, Larijani B. Nutrition and diet therapy in diabetes mellitus: A roadmap based on available evidence. *J Diabetes Metabolic Dis*. 2021;20(2):1913-8. DOI: <https://doi.org/10.1007/s40200-021-00876-2>
 40. Kanaley JA, Colberg SR, Corcoran MH, Malin SK, Rodriguez NR, Crespo C J et al. Exercise/physical activity in individuals with type 2 diabetes: A consensus statement from the American College of Sports Medicine. *Med Sci Sports Exerc*. 2022;54(2):353-68. DOI: <https://doi.org/10.1249/MSS.0000000000002800>
 41. Neves Júnior TT, Rêgo-de Queiroz AA, Araújo CE, Assis SCJ, Nóbrega ETM, Perfil clínico y sociodemográfico de usuarios con enfermedades crónicas en atención primaria de salud. *Enferm. glob. [Internet]*. 2023 [citado 21 jun. 2023];22(69):245-82. DOI: <https://doi.org/10.6018/eglobal.512211>
 42. Bohnert KL, Zellers JA, Jeong H J, Chen L, York A, Hastings M K. Remote research: Resources, intervention needs, and methods in people with diabetes and peripheral neuropathy. *Journal of Diabetes Science and Technology*. 2023;17(1):52-8. DOI: <https://doi.org/10.1177/19322968221103610>
 43. Kokoszka A. Treatment adherence in patients with type 2 diabetes mellitus correlates with different coping styles, low perception of self-influence on disease, and depressive symp-

- toms. Patient Preference and Adherence. 2017;11:587-95. DOI: <https://doi.org/10.2147/PPA.S124605>
44. Mahdizadeh M, Peyman N, Thagipour A, Esmaily H. Effect of health education program on promoting physical activity among diabetic women in Mashhad, Iran: Applying social cognitive theory. *J Res Health Sci.* 2013;13(1):90-7.
45. Carvalho LK da CAA, Ibiapina TF, Rodrigues M da CVA, Leite MAE, Lopes de CM et al. Capacitação de enfermeiros na Estratégia Saúde da Família: análise do processo de educação permanente para o Sistema Único de Saúde. *Nursing.* 2018;21(247):2506-12. DOI: <https://doi.org/10.36489/nursing.2018v21i247p2506-2512>
46. Giovanela L, Mendonça MHM de, Buss PM, Gadelha CAG, Galvão LAC et al. De Alma-Ata a Astana. Atenção primária à saúde e sistemas universais de saúde: compromisso indissociável e direito humano fundamental. *Cad Saúde Pública* 2019;35(3):e00012219. DOI: <https://doi.org/10.1590/0102-311X00012219>
47. Pereira VS, Souza CA de, Louro TQ, Oliveira ES de, Lima DM, Silva RCL da, Marta CB. Avaliação de Tecnologias em Saúde: Estado da Arte. *Saud Coletiv.* 2019;9(51):2035-40. DOI: <https://doi.org/10.36489/saudecoletiva.2019v9i51p2035-2040>
48. Teston EF, Spigolon DN, Maran E, Santos A de L, Matsuda LM et al. Perspectiva de enfermeiros sobre educação para a saúde no cuidado com o Diabetes Mellitus. *Rev Bras Enferm.* 2018;71(supl. 6):2899-907. DOI: <https://doi.org/10.1590/0034-7167-2018-0396>
49. Rawshani A, Rawshani A, Franzén S, Sattar N, Eliasson B et al. Risk factors, mortality, and cardiovascular outcomes in patients with Type 2 Diabetes. *N Engl J Med.* 2018;379(7):633-44. DOI: <https://doi.org/10.1056/NEJMoa1800256>
50. Cesarino CB, Borges PR, Ribeiro R de CHM, Ribeiro DF, Kusumota L. Avaliação do risco cardiovascular de pacientes renais crônicos segundo critérios de Framingham. *Acta Paul Enferm.* 2013;26(1):101-7. DOI: <https://doi.org/10.1590/S0103-21002013000100016>

Appendix A. Data Extraction Form

STUDY CHARACTERISTICS	
Authors	
Title	
Publication year	
Study country	
Journal name	
Document type	
Language	
Study objective	
Study scenario	
Methods	
Population/sample size within the evidence source (if applicable)	
Inclusion criteria	
Exclusion criteria	
Data collection instruments/techniques	
Results (program/intervention impact)	
Conclusions/key aspects that relate to the research question	
PROGRAM CHARACTERISTICS	
PARTICIPANTS	
Social (age, housing, marital status)	
Clinical	
INTERVENTION/PROGRAM CHARACTERISTICS	
Program/intervention duration	
Baseline theoretical model	
Resources employed (material/ pedagogical, and human)	
Program/intervention implementation	
PROGRAM RESULTS	
Program impact	
User satisfaction evaluation	

OSF REGISTRIES
Add New My Registrations Help Donate Join Login

NURSE-LEADED PROGRAMS FOCUSING ON SOCIAL SUPPORT FOR PEOPLE WITH TYPE 2 DIABETES: SCOPING REVIEW

Public registration
Updates

Overview

- Files
- Resources
- Wiki
- Components 0
- Links 0
- Analytics
- Comments 0

Open practice resources

- Data
- Analytic code
- Materials
- Papers
- Supplements

Study Information

Hypotheses

What nurse-led programs exist to support the self-management of people with type 2 diabetes through social support and what are their characteristics?

In order to answer the question mentioned above, a preliminary exploratory search was carried out in the Worldwide Science, Stelo, Medicine (via PubMed), and CINAHL (via EBSCO HOST) databases, JBI Evidence Synthesis, and the lack of systematic review was observed, or scope analysis, current or in progress, on the aforementioned topic.

Thus, it can be seen that there is a research gap urging me to carry out a scoping review to map the theme in the literature.

Design Plan

Study type
Other

Blinding
No blinding is involved in this study.

Is there any additional blinding in this study?
No response

Study design
A favorable situation regarding the benefits of monitoring the chronic disease with engaged participation from the family, effective to the professional during the treatment, that is, involving emotional and educational support, has shown significant impacts, since the patient is able to acquire courage and witpower in the assiduous continuity of their self-care and health management (JONQUIM, 2022).

Another factor that impacts adherence and continuity of the therapeutic regimen is interconnected with the appropriate form of communication for the recipient. For this to occur properly, professionals must be attentive and qualified since educational management is a very important basis within social support (OLIVEIRA LIMA, 2017).

In addition, the educational aspects provided by professionals trained in the composition of care for patients with type 2 diabetes mellitus contribute to the health-disease process in a way that empowers the client. Therefore, it is necessary to carry out this Scoping Review with the proposal to unite and synthesize current scientific evidence on the subject.

No files selected

Randomization
No response

Sampling Plan

Existing data
Registration prior to analysis of the data

Explanation of existing data
No response

Data collection procedures
The review will be conducted following the Joanna Briggs Institute (JBI) methodological guidelines.

Based on the PCC strategy, with "P" equivalent to the study population, "C" for the concept to be investigated, "C" for the context to be addressed. Therefore, the population is adults who have type 2 DM; the concept, nurse-led social support programs; the context, in primary health care or in the community.

The Boolean phrase ("Self Care"[MeSH Terms] OR "Self Care"[Text Word] OR "Self management"[Text Word]) AND ("Social Support"[MeSH Terms] OR "Social Care"[Text Word] OR "Online Social Support"[Text Word] OR "Perceived Social Support"[Text Word] OR "Perceived Social Supports"[Text Word]) AND ("nurse-led education"[Text Word] OR "nurse-delivered intervention"[All Fields] OR "nurse-led intervention" OR "program development"[All Fields] OR "nurse-led program"[All Fields]) AND ("diabetes" OR "diabetes mellitus" OR "diabetes mellitus, Type 2" OR "Diabetes Mellitus, Type II" OR "Type 2 Diabetes Mellitus" OR "Type 2 Diabetes" OR "Diabetes, Type 2"), built with the technical support of a USF librarian, it will serve as a guide for searches in the databases chosen to be part of the study.

Research will be carried out in the following databases: Web of Science, Medicine (PubMed), CINAHL (EBSCO), Catalog of Theses and Dissertations = Capes, Scopus Preview, JBI Evidence Synthesis, PsycInfo (APA), Open Access Scientific Repositories of Portugal.

No files selected

Sample size
There will be a prior search in the literature.

Sample size rationale
No response

Stopping rule
No response

Variables

Manipulated variables
No response
No files selected

Measured variables
It is a scoping review.
No files selected

Indices
No response
No files selected

Analysis Plan

Statistical models
Data analysis will take place after study selection with reanalysis of criteria present in the study form. The studies defined to be part of the scoping review will undergo a filing, consisting of emphasis on content.

The data will be presented through narrative summaries and, if necessary, flowcharts and tables accompanied by texts, regardless of whether they are self-explanatory.

No files selected

Transformations
No response

Inference criteria
No response

Data exclusion
No response

Missing data
No response

Exploratory analysis
No response

Other
No response

Contributors
Isabela Mendonça Rodrigues dos Santos, Geysa Santos Góis Lopes, and Isaura Leticia Tavares Paimeira Rolim

Description
Programs led by nurses to support self-management of care from the perspective of social support, in adults with type 2 diabetes.

Registration type
OSF Preregistration

Date registered
November 6, 2022

Date created
November 6, 2022

Associated project
osf.io/j3ymk

Internet Archive link
<https://archive.org/details/osf-registrations-4wv6-v1>

Category
Project

Registration DOI
<https://doi.org/10.17605/OSF.IO/WVWT6>

Subjects

- Life Sciences
- Medicine and Health Sciences

License
Academic Free License (AFL) 3.0

Citation
osf.io/4wv6