

Sección Agrícola

Digestion of the inhibitor α AI by *Hypothenemus hampei* aspartic proteinases

Digestión de un inhibidor de α AI por aspartico proteinasas de *Hypothenemus hampei*

ARNUBIO VALENCIA J.^{1*}, JORGE W. ARBOLEDA V.²

Summary. The digestive fluid of the coffee berry borer, *Hypothenemus hampei*, larvae contains aspartic proteases, which show greatest activity at pH 3.0 and a temperature of 40°C. By using isoelectric focusing gels, it was possible to observe two well-defined bands in the intestinal tract. The band of highest activity was found very close to the anodic side and showed a *pI* of 5.2. The addition of 0.1 μ M of Pepstatin A caused a 70% inhibition of protease activity and a 90% inhibition when the concentration was 1 μ M. The strong inhibition of the protease activity in the intestinal tract of *H. hampei* after addition of Pepstatin A, allowed us to identify this protease as an aspartic-type protease. Other results permitted us to classify this protease as Cathepsin D given that when BSA was used as a substrate the enzyme only showed 20% of the total activity with respect to hemoglobin. Incubation of the α AI inhibitor from *Phaseolus vulgaris* with increasing amounts of this intestinal enzyme from the coffee berry borer, showed that the biological activity of the inhibitor was significantly reduced due to the proteolytic activity. This work constitutes the first electrophoretic evidence of α AI proteolysis of *P. vulgaris* by endogenous proteases from *H. hampei*.

Key words: Enzymes. Coffee berry borer. Cathepsin.

Resumen. El fluido digestivo de las larvas de la broca de café, *Hypothenemus hampei* contiene aspártico proteasas, las cuales muestran una gran actividad a pH 3.0 y a una temperatura de 40°C. Mediante el uso de gels de isoelectroenfoque (IEF) fue posible observar dos bandas de actividad bien definidas en el tracto intestinal. La banda de mayor actividad que se encontró muy cerca del ánodo presentó un *pI* de 5.2. La adición de pepstatin A, a una concentración de 0.1 μ M causó una inhibición del 70% de la actividad proteasa y de un 90% cuando la concentración del inhibidor fue de 1 μ M. La fuerte inhibición de la actividad proteasa presente en el tracto intestinal de *H. hampei*, después de la adición del pepstatin A, permitió identificar esta proteasa como del tipo aspártico. Otros resultados permitieron clasificar esta enzima como Cathepsin D, dado que cuando se empleó BSA como sustrato la enzima mostró solo un 20% del total de actividad con respecto a hemoglobina. La incubación del inhibidor de α -amilasas (α AI) de *Phaseolus vulgaris*, con cantidades crecientes de esta enzima intestinal de la broca de café, mostró que la actividad biológica del inhibidor fue reducida significativamente debido a la actividad proteolítica. Este trabajo se constituye en la primera evidencia electroforética de la proteólisis del α AI de *P. vulgaris* por proteasas endógenas de *H. hampei*.

Palabras clave: Enzimas. Broca del café. Cathepsin.

Introduction

The coffee berry borer, *Hypothenemus hampei* (Ferrari) (Coleoptera: Curculionidae: Scolytinae) is not only a major pest in Colombia and other countries, but also a serious economic pest of coffee in many coffee-growing areas of the world (Murphy and Moore 1990; Reid 1983). Insects, as well as other organisms, use proteinase for digestion of proteins that are ingested with their food (Xu and Qin 1994). Usually, the proteinase type is determined by the pH range of activity, by its similarity to well-characterized proteinases and by its sensitivity or susceptibility to different inhibitors (North 1982). Several insect proteases have already been described, isolated and characterized, some of which occur as mixtures of different isozymes, and which are generally simi-

lar in their catalytic activity to those found in vertebrates (Blanco *et al.* 1996; Applebaum 1985). However, some proteinases present major differences in their more important characteristics such as: location, optimal pH activity, kinetic constants and thermostability (Blanco *et al.* 1996).

Most of the midgut proteolytic enzymes in coleopteran insects have been shown to be cysteine proteinases (Murdock *et al.* 1987), or aspartic proteinases often found in combination with other proteinases and which are very active in acid environments, pH normally below 4.5 (Jongsma and Bolter 1997).

Characterizing the major protease activity present in the digestive tract of *H. hampei* larvae is important not only to

understand the protein digestion, but also to know how susceptible to digestion the α AI from *Phaseolus vulgaris* is. Interest in this area has been stimulated by the finding that insect digestive proteases have potential to digest α Amylase inhibitors and to increase by this way their natural resistance to these plant inhibitors (Ishimoto and Chrispeels 1996).

This paper describes the use of specific and general substrates, with a specific inhibitor, to determine the type of protease in the intestinal tract of larvae of *H. hampei* and to assess the role of this protease in the α AI digestion. We believe the present paper to be not only the first report but also the first electrophoretic evidence by using zymograms of α AI proteolysis with endogenous aspartic proteases from *H. hampei*.

1* Corresponding author: Departamento de Fitotecnia. Facultad de Ciencias Agropecuarias. Universidad de Caldas. Calle 65 # 26-10. Manizales-Colombia. Tel. + 57-68-812975. Fax: + 57-68-812975. E-mail: arnubio@laciudad.com

2 Disciplina de Entomología. Centro Nacional de Investigaciones de Café, CENICAFÉ. Colombia. Tel.: +57-68-506550. A.A. 2427 Manizales. E-mail: jwarboleda@terra.com.co

Materials and Methods

Sources of plant inhibitors and insects

Hypothenemus hampei was obtained locally in Colombia and cultivated in an insectary at Cenicafe in Chinchiná, Colombia. *P. vulgaris* cv. L. radical is a local commercial bean in Manizales, Colombia.

Extraction procedures of insect proteases

Fifty insect guts carefully obtained by dissection of whole coffee berry borer larvae were homogenized in 50 µl of citrate buffer 0,05M pH 3,0, filtrated and centrifuged at 10.000g for 30 min at 4°C. The supernatant was freeze-dried and the resulting powder was stored at -80°C and used as a source of enzyme.

Purification of αAI-1

Seeds (20 g) of *P. vulgaris* were ground in a coffee grinder and the powder mixed with 250 ml of distilled water and stirred in the cold room from 6h. The slurry was centrifuged for 30 min at 8000g and the pH of the supernatant was adjusted to 4,0 with 1 N HCl. The precipitated proteins were removed by centrifugation (30 min at 8000g) and the pH of the supernatant adjusted to pH 6,7 with 1 M KPO₄. If there was a precipitate it was removed again by centrifugation. The proteins in the clear supernatant were precipitated by the gradual addition of saturated ammonium sulfate (4,1M) to a final concentration of 85% saturation.

After standing on ice for 4 h, the proteins were collected by centrifugation for 30 min at 8000g. The pellet was resuspended in water, dialyzed against water for 36 h and then against 20 mM KPO₄ pH 6,7 for at least 12 h. The protein solution was loaded on a DEAE cellulose column (120 x 16 mm) and the column washed with 200 ml of the same phosphate buffer. The column was eluted with a linear NaCl gradient (0 to 100 mM) and absorbance at 280 nm and αAI activity were measured in the eluted fractions.

Protein determination

Protein was determined by the Coomassie procedure (Bradford 1976). Bovine serum albumin was used as standard protein.

Electrophoretic separation and enzyme zymogram in isoelectrofocusing (IEF gels)

For all gels we used the PhastSystem electrophoresis unit (Pharmacia) following the manufacturer instructions. The isoelectric

point was calculated by using standard proteins with known isoelectric points and with software (ImageMaster VDS) provided by Pharmacia Biotech. For the protease zymogram, the proteins were first separated in nondenaturing conditions on IEF (3,0-9,0) Phastgels. After electrophoresis, the IEF gel was lightly rinsed with distilled water and incubated for 3 hours at 35°C with Citrate buffer 0,05M pH 3,0 containing 1% of hemoglobin. The gel was rinsed with water and stained with Coomassie Blue (Electrophoresis solution) during 30 min at 45°C and cleaned with water / methanol / acetic acid (60:30:10) solution. Protease activity appears on the polyacrylamide gel as clear bands on a blue-colored background. The gel was washed once again with water and then photographed.

Electrophoretic separation and inhibitor zymogram in isoelectrofocusing (IEF)

For the inhibitor zymograms, the IEF gels were incubated in 1,5% soluble starch in 10 mM sodium citrate, pH 5,0, containing 10 mM NaCl and 20 mM CaCl₂ for 1h at 4°C. The starch-gel was rinsed with water and incubated for 20 min at 30°C, rinsed and incubated with human salivary amylase dissolved in the same buffer. Staining with KI/I₂ (I: 0,5% and KI: 5%) produced dark blue bands on a light background (Valencia *et al.* 2000).

Proteolytic activity

Total proteolytic activity was assayed according to the method of Lenney (1975) and Blanco Labra *et al.* (1996). The bovine hemoglobin was prepared in water and denaturalized at pH 1.0 with a HCl solution 6,0N. This solution was dissolved with Citrate buffer pH 3,0 in order to get a 0,083% hemoglobin final solution. To carry out the activity tests, 40 µl of the enzyme solution was incubated for 3 hours at 30°C with 300 µl of Citrate buffer (0,05M pH 3,0) and 1,3 ml of hemoglobin solution (0,083%). The reaction was stopped by the addition of 2,5 ml of Trichloroacetic acid solution 0,34N. Then, samples were put in ice for 10 minutes and centrifuged at 10000g for 10 minutes. One enzymatic activity unit was defined as a decrement of 0,001 in absorbance at 280 nm. The absorbance was determined by using a UV/VIS spectrophotometer (UNICAM).

Proteolytic inhibition by pepstatin

Proteolytic inhibition was evaluated following preincubation of the enzyme

preparation with pepstatin A, a synthetic inhibitor. This specific inhibitor was added to enzyme preparation at a final concentration of 0,1 µM, 0,4 µM, and 1 µM and allowed to stand at 25°C for 15 min before adding the substrate. Following preincubation the remaining activity was assayed as described above.

Proteolytic digestion of αAI

Proteolytic digestion of αAI (*Phaseolus vulgaris*) by acidic proteases present in the digestive fluid of *H. hampei*, was assayed as follows: An aliquot (1 µl) of the pure αAI (2 µg/µl) was incubated for two hours at 30°C with 1 µl of crude enzyme extract containing 3,92, 1,96, 0,98 or 0,0 enzymatic activity units. The reaction was stopped by adding Pepstatin A to this reaction solution at a final concentration of 0,1 µM. 1 µl of this solution containing the αAI was separated by using an isoelectrofocusing gel (IEF 3-9). Inhibitor zymogram was developing as described above.

Results and Discussion

Characteristics of intestinal tract proteases

In preliminary experiments we determined some biochemical properties of the digestive proteases of the coffee berry borer larvae. We found, that the enzyme has an optimum pH between 2,5 and 3,5 and it holds an 80% of their enzymatic activity when it is heated up to 50°C. In addition, when different buffer solutions were assayed, it was found that 0,05 M Citrate buffer, pH 3,0, was the best buffer solution to assay the proteolytic activity (Data not shown). Similar to most insect proteases, the enzyme activity found in coffee berry borer has similar values to those registered previously for other Coleoptera insects. For example, *Tribolium castaneum* (Herbst) shows an optimum proteolytic activity at pH between 3,0 and 6,9 (Blanco *et al.* 1996; Murdock *et al.* 1984; Birk *et al.* 1962); between 3,3 and 6,0 for *Callosobruchus maculatus* (F.) (Silva y Filho 1991; Murdock *et al.* 1984; Kitch y Murdock 1986; Gatehouse *et al.* 1985; Campos *et al.* 1989); between 5,5 and 6,5 for *Acanthoscelides obtectus* Say (Wieman y Nielsen 1988); and between pH 3,5 and 5,5 for *Zabrotes subfasciatus* (Boh) (Lemos *et al.* 1990). The highest proteolytic activity founded at pH 3,0 in *H. hampei* clearly indicates that this enzyme is an acid protease (Aspartic protease).

Similar results have been reported in other Coleoptera insects (Vundla *et al.* 1992; Barret 1977; Houseman and Downe 1982; Okasha 1968; Smith and Birt 1971).

The gradual loss of proteolytic activity that was observed in *H. hampei* over 50°C, has also been registered for proteinases coming from other organisms such as *Mujil auratus* (Bonete *et al.* 1984), *Tribolium castaneum* (Blanco *et al.* 1996), and *Trichoplusia ni* (Pritchett *et al.* 1981), indicating its thermal instability.

Electrophoretic zymograms and *pI* determination

To find out if coffee berry borer insect can synthesize different isoforms of acidic proteases, we analyzed enzymatic extracts from its intestinal tracts on native 16% polyacrylamide gels and develop them as zymograms. We observed two well-defined bands in the intestinal tract of *H. hampei* larvae (Fig. 1).




Figure 1. Zymogram of proteolytic activity from the digestive tract of *H. hampei* by using native electrophoresis on 7,5% homogenous Phastgel. After electrophoresis the gel was incubated with 1% hemoglobin solution in buffer Citrate pH 3,0, during 2 h at 35°C.

Similar analysis of the acidic proteases after separation of the proteins present in the intestinal tract on an isoelectric-focusing gels (IEF 3-9) showed the presence of a single major band very close to the anodic side with a *pI* of 5,2 (Fig. 2). Again, this result with *H. hampei* proteases was in general agreement with those reported for other insect acidic proteases, most of which fall in the acidic region of the IEF gel. For example, the two-reported aspartic proteases of *Tribolium castaneum* had isoelectric points between 2,8 and 3,5, respectively (Blanco *et al.* 1996; Barret 1977). The presence of this enzyme in the intestinal lumen of the insect shows the digestive nature of the enzyme.




Figure 2. Determination of the *pI* of the major aspartic protease from intestinal tract of *H. hampei* by using an IEF Phastgel (3-9). Lane 1, represents the major aspartic protease *H. hampei* larvae. Lane 2, displays the *pI* markers.

Enzymatic inhibition by pepstatin A

Pepstatin A is a potent inhibitor of aspartic proteases and we prove its effect on *H. hampei* acidic proteases. So, to determine if the enzyme activity that is present in the digestive tract of coffee berry borer, could be inhibited by Pepstatin A, we assayed an intestinal extract for protease activity after incubation with increasing amounts of Pepstatin A. The results (Fig. 3) showed that addition of 0,1 μ M of Pepstatin A caused a 72% inhibition of protease activity and a 91% inhibition when 1 μ M of inhibitor was assayed. The strong inhibition caused on the protease activity in *H. hampei* after addition of Pepstatin A, allow us to conclude that the endogenous proteolytic activity in the intestinal tract of coffee berry borer is aspartic protease type.

In order to determine if the insect protease was Pepsin or Cathepsin D type, we assayed the enzyme activity by using hemoglobin and bovine serum albumin (BSA) as two different substrates. The results showed that this proteolytic enzyme hydrolyze hemoglobin more efficiently than BSA. Only 20% of the total activity on hemoglobin was found when BSA was used as substrate (data not shown).

According to this, it was possible to classify this enzyme as a protease cathepsin D type. The results of our study are consistent with the identification of a cathepsin protease activity in intestinal tracts of *Tribolium* species (Blanco *et al.* 1996).




Figure 3. Effect of Pepstatin A on aspartic protease activity of coffee berry borer (*H. hampei*) gut extract. Data are presented as percentage of control containing no inhibitor.




Figure 4. Proteolytic digestion of α AI-1 (*Phaseolus vulgaris* c.v. Radical) by *H. hampei* larval acidic protease(s). Purified α AI-1 was incubated with increasing amounts of larval midgut acidic protease from Coffee berry borer. Lane 1, 3,92 units/ μ l. Lane 2, 1,96 units/ μ l. Lane 3, 0,98 units/ μ l. Lane 4, Control (Untreated α AI-1). An aliquot (1 μ l) of the digestion solution containing the remaining α AI-1 (1 μ g/ μ l) was loaded in each well and run on an IEF (3-9) gel.

Proteolysis of α AI by larval proteases

To determine if the α AI-1 could be digested by the acidic proteases found in the intestinal tract of coffee berry borer, we assayed an exact quantity of α AI-1 for inhibitory activity after incubation with increasing amounts of acidic protease activity. The results (Fig. 4) showed that the inhibitory activity of α AI was significantly reduced when the inhibitor was incubated with acidic proteases from the intestinal tract of *H. hampei*. 3,92 units/ μ l of aspartic proteases from coffee berry borer nearly destroyed the total biological activity of the α AI from *P. vulgaris* after two hours of incubation. Previous works of proteolysis of α AI by larval enzymes, showed that *Z. subfasciatus* larval enzymes can cleave α AI-1 to smaller molecules (Ishimoto and Kitamura 1992). Ishimoto and Chrispeels (1996) have reported a similar result when seed extracts of *Phaseolus vulgaris* were incubated with larval midgut extracts of *Z. subfasciatus*, *Acanthoscelides obtectus*, and *Callosobruchus chinensis*, and the resulting α AI polypeptides were visualized on an immunoblot after SDS-PAGE. Although it is well known that insect gut proteases could digest different proteins that are intake by the insect during feeding, it doesn't exist electrophoretic evidence on this phenomenon when the well-characterized α AI is used as target; especially if the biological activity of the amylase inhibitor is monitored.

Because of this evidence, use of transgenic insect-resistant plant expressing α AI requires not only a complete characterization of the digestive proteases in the insect gut but also, requires to know if the inhibitor can act at the pH found in

its digestive tract and, if it is resistant to attack by insect gut proteases.

Based on the experiments presented here, we think that in order to increase the protective efficacy and durability of α AI in transgenic plants, it is envisaged that "Packages" of different genes must be introduced into plants. The component of such packages should each act on different targets within the insect. Protease inhibitors should be particularly valuable especially because they would protect other introduced gene products from premature and active digestion in the insect gut.

Conclusions

- The results in this paper provide further insight into the relation between α -amylase inhibitors and the digestive proteases that are present in the insect target.
- The data presented here lead to three important conclusions. First, the endogenous proteolytic activity in the digestive tract of *H. hampei* is aspartic protease (Cathepsin D) type. Second, the inhibitory activity of α AI could be significantly reduced when the inhibitor is incubated with acidic proteases from the intestinal tract of coffee berry borer. Third, protease inhibitors should be particularly valuable especially because they would protect other introduced gene products from premature and active digestion in the insect intestinal tract.

Acknowledgements

The authors gratefully acknowledge Colciencias and Cenicafé (Colombia) for financial support. We also thank the Entomology Laboratory of Cenicafé for their assistance with the insects.

Literature cited

- APPLEBAUM, S. W. 1985. Biochemistry of digestion. *Comprehensive Insect Physiology Biochemistry and Pharmacology* 4: 279-311.
- BARRET, A. J. 1977. Cathepsin D and other carboxyl proteinases from mammalian cells and tissues. Amsterdam, North-Holland. p. 209-248.
- BIRK, Y.; HARPAZ, I.; ISHAAYA, I.; BONDY, A. 1962. Studies on the proteolytic activity of the beetles *Tenebrio* and *Tribolium*. *Journal of Insect Physiology* 8: 417-429.
- BLANCO, L. A.; MARTÍNEZ, N. A.; SANDOVAL, L.; DELANO, J. 1996. Purification and characterization of a digestive cathepsin D proteinase isolated from *Tribolium castaneum* larvae (Herbst). *Insect Biochemistry and Molecular Biology* 26(1): 95-100.
- BONETE, M. J.; MANJON, A.; LLORCA, F.; IBORRA, J. Z. 1984. Acid proteinase activity in fishes II. Purification and characterization of Cathepsin B and D from *Mujil auratus* muscle. *Comparative Biochemistry and Physiology* 78B: 207-213.
- BRADFORD, M. M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry* 72: 248-254.
- CAMPOS, F. A. P.; XAVIER-FILHO, J.; SILVA, C. P.; ARY, M. B. 1989. Resolution and partial characterization of proteinases and α -amylases from midguts of larvae of the bruchid beetle *Callosobruchus maculatus* (F.). *Comparative Biochemistry and Physiology* 92B: 51-57.
- GATEHOUSE, A. M. R.; BUTLER, K. J.; FENTON, K. A.; GATEHOUSE, J. A. 1985. Presence and partial characterization of a major proteolytic enzyme in the larval gut of *Callosobruchus maculatus*. *Entomologia Experimentalis et Applicata* 39: 279-286.
- HOUSEMAN, J. G.; DOWNE, A. E. R. 1982. Characterization of an acidic proteinase from the posterior midgut of *Rhodnius prolixus* Stål (Hemiptera: Reduviidae). *Insect Biochemistry* 12(6): 651-655.
- ISHIMOTO, M.; CHRISPPEELS, M. J. 1996. Protective mechanism of the Mexican Bean Weevil against high levels of α -amylase inhibitor in the common bean. *Plant Physiology* 111: 393-401.
- ISHIMOTO, M.; KITAMURA, K. 1992. Tolerance to the seed α -amylase inhibitor by the two insect pests of the common bean, *Zabrotes subfasciatus* and *Acanthoscelides obtectus* (Coleoptera: Bruchidae). *Applied Entomology and Zoology* 27: 243-251.
- JONGSMA, M. A.; BOLTER, C. 1997. The adaptation of insect to plant protease inhibitors. *Journal of Insect Physiology* 43(10): 885-895.
- KITCH, L. W.; MURDOCK, L. L. 1986. Partial characterization of a major midgut tliol proteinase from larvae of *Callosobruchus maculatus* (F.). *Archives of Insect Biochemistry and Physiology* 3: 561-576.
- LEMONS, F. J.; CAMPOS, F. A. P.; SILVA, C. P.; XAVIER-FILHO, J. 1990. Proteinases and amylases of larval midgut of *Zabrotes subfasciatus* reared on cowpea (*Vigna unguiculata*) seeds. *Entomologia Experimentalis et Applicata* 56: 219-227.
- LENNEY, J. F. 1975. Three yeast proteins that specifically inhibit yeast proteases A, B and C. *Journal of Bacteriology* 122: 1265-1273.

- MURDOCK, L. L.; BROOKHART, G.; DUNN, P. E.; FOARD, D. E.; KELLY, S.; KITCH, L.; SHADE, R. E.; SHUCKLE, R. H.; WOLFSON, J. L. 1984. Cysteine digestive proteinases in Coleoptera. *Comparative Biochemistry and Physiology* 87B: 783-787.
- MURDOCK, L. L.; BROOKHART, G.; DUNN, P. E.; FOARD, D. E.; KELLEY, S.; KITCH, L.; SHADE, R. E.; SHUCKLE, R. H.; WOLFSON, J. L. 1987. Cysteine digestive proteinases in Coleoptera. *Comparative Biochemistry and Physiology* 87: 783-787.
- MURPHY, S. T.; MOORE, D. 1990. Biological control of the coffee berry borer *Hypothenemus hampei* (Ferrari) (Coleoptera: Scolytidae), previous programs and possibilities for the future. *Biocontrol News and Information* 11: 107-117.
- NORTH, M. J. 1982. Comparative biochemistry of the proteinases of eucaryotic microorganisms. *Microbiology* 46: 308-340.
- OKASHA, A. Y. K. 1968. Effect of sub-lethal temperature on an insect, *Rhodnius prolixus* (Stal) III. Metabolic changes and their bearing on the cessation and delay of moulting. *Journal of Experimental Biology* 48: 475-486.
- PRITCHETT, D. W.; YOUNG, S. Y.; GEREN, C. R. 1981. Proteolytic activity in the digestive fluid of larvae of *Trichoplusia ni*. *Insect Biochemistry* 11(5): 523-526.
- REID, J. C. 1983. Distribution of the coffee berry borer *Hypothenemus hampei*, within Jamaica, following its discovery in 1978. *Tropical pest management* 29: 224-230.
- SILVA, C. P.; XAVIER-FILHO, J. 1991. Comparison between the levels of aspartic and Cysteine proteinases of *Callosobruchus maculatus* (F) and *Zabrotes subfasciatus* (Boh) (Coleoptera: Bruchidae). *Comparative Biochemistry and Physiology* 99B: 529-533.
- SMITH, E.; BIRT, L. M. 1971. Proteolytic activity during the metamorphosis of the blowfly, *Lucila*. *Insect Biochemistry* 2: 218-225.
- VALENCIA, J. A.; A. E. BUSTILLO, G. A. OSSA.; CHRISPPEELS, M. J. 2000. Amylases of the coffee berry borer (*Hypothenemus hampei*) and their inhibition by two plant amylase inhibitors. *Insect Biochemistry and Molecular Biology* 30: 207-213.
- VUNDLA, W. R. M.; BROSSARD, M.; PEARSON, D. J.; LABONGO, V. L. 1992. Characterization of aspartic proteinases from the gut of the Tick, *Rhipicephalus appendiculatus* Neuman. *Insect Biochemistry Molecular Biology* 22(4): 405-410.
- WIEMAN, K. F.; NIELSEN, S. S. 1988. Isolation and partial characterization of a major gut proteinase from larval *Acanthoscelides obtectus* Say (Coleoptera: Bruchidae). *Comparative Biochemistry and Physiology* 89B: 419-426.
- XU, G.; QIN, J. 1994. Extraction and characterization of midgut proteases from *Heliothis armigera* and *H. assulta* (Lepidoptera: Noctuidae) and their Inhibition by tannic acid. *Journal of Economic Entomology* 87(2): 334-338.

Recibido: 12-oct-04 • Aceptado: 23-dic-04

