

Artículo original

Violence Exposure and Disability in Colombian Female Survivors of Intimate Partner Violence: the Mediating Role of Depressive Symptoms



Lina Marcela Sandoval^{a,b,*}, Elvis Siprian Castro^c, Andrés Fandiño-Losada^a, Sara Gabriela Pacichana-Quinayaz^a, Shirley Evelyn Lennon^a, María Isabel Gutiérrez-Martínez^a

^a School of Health, Cisalva Institute, Universidad del Valle, Cali, Colombia

^b Institución Universitaria Escuela Nacional del Deporte, Cali, Colombia

^c School of Human Rehabilitation, Grupo SINERGIA, Universidad del Valle, Cali, Colombia

ARTICLE INFO

Article history:

Received 15 August 2021

Accepted 21 November 2021

Available online 19 January 2022

Keywords:

Disability

Intimate partner violence

Depressive symptoms

Mediating

Path analysis

ABSTRACT

Background: Violence is an important public health problem and one of the main causes of deaths worldwide. The mental health consequences of surviving intimate partner violence (IPV) include depression, anxiety and post-traumatic stress disorder. Previous studies have identified that there is a relationship between depression and level of disability in female survivors of IPV. Estimating the direct, indirect or total effect of an exposure on an outcome makes it possible to identify mediating effects between a group of variables. Detecting mediation effects is useful for identifying casual pathways that generate a final outcome and provides a rationale for designing interventions to target the mediator, which in turn positively affects the outcome. The objective was to identify the mediating role of depressive symptoms on the relationship between IPV and disability.

Methods: This was a cross-sectional study of 94 women over the age of 18 who were survivors of IPV by men. They were recruited from two public hospitals in Cali and Tuluá in southwest Colombia. An analysis of casual relationships was performed using structural equation modelling that was made up of: four exogenous observed variables (age, current relationship status [in a relationship or single], level of schooling, and history of an impairment), intermediate endogenous variables (violence and depressive symptoms), and the main endogenous variable (disability). The analyses were carried out in Stata14.2.

Results: The direct effect of IPV severity on the level of disability was not statistically significant ($\beta = 0.09$; $P = 0.63$). However, the indirect effect of IPV severity on disability mediated by depressive symptoms was ($\beta = 0.39$; $P < 0.01$). The total effect of IPV severity on the level of disability was even greater ($\beta = 0.48$; $P = 0.01$).

* Corresponding author.

E-mail address: lina.sandoval@correounivalle.edu.co (L.M. Sandoval).

<https://doi.org/10.1016/j.rcp.2021.11.009>

0034-7450/© 2021 Asociación Colombiana de Psiquiatría. Published by Elsevier España, S.L.U. All rights reserved.

Conclusions: This study found a complete mediating role of depressive symptoms on the relationship between the severity of IPV and the level of disability for the female participants in this study. The results of this research contribute to defining strategies to prevent and address intimate partner violence, depressive symptoms and disability in this population.

© 2021 Asociación Colombiana de Psiquiatría. Published by Elsevier España, S.L.U. All rights reserved.

Exposición a la violencia y discapacidad en mujeres colombianas sobrevivientes de violencia de pareja íntima: el papel mediador de los síntomas depresivos

R E S U M E N

Palabras clave:

Discapacidad
Violencia íntima de pareja
Síntomas depresivos
Mediación
Análisis de ruta

Introducción: La violencia es un importante problema de salud pública y una de las principales causas de muerte en todo el mundo. Las consecuencias para la salud mental de sobrevivir a la violencia de la pareja íntima incluyen depresión, ansiedad y trastorno de estrés postraumático. Estudios previos han identificado que existe una relación entre la depresión y el nivel de discapacidad en mujeres sobrevivientes a violencia de pareja íntima. La estimación del efecto directo, indirecto o total de una exposición en un resultado permite identificar efectos mediadores en un grupo de variables. La detección de los efectos de la mediación es útil para identificar vías causales que generan un resultado final y proporciona una justificación para diseñar intervenciones dirigidas al mediador, lo que a su vez afecta positivamente al resultado. El objetivo es identificar el efecto mediador de los síntomas de depresión en la relación entre gravedad de la violencia y el nivel de discapacidad de las mujeres víctimas de violencia de pareja.

Métodos: Estudio transversal que incluyó a 94 mujeres de 18 o más años víctimas de violencia de pareja, identificadas en 2 instituciones públicas de salud de Cali y Tuluá, en el suroccidente de Colombia. Se realizó un análisis de rutas causales mediante un modelo de ecuaciones estructurales (SEM), en Stata[®] 14.

Resultados: El efecto directo de la gravedad de la violencia en la discapacidad no es significativo ($\beta = 0,09$; $p = 0,63$); sin embargo, el efecto indirecto mediado por los síntomas de depresión incrementa su magnitud y es significativo ($\beta = 0,39$; $p < 0,01$). El efecto total es mayor y significativo ($\beta = 0,48$; $p = 0,01$).

Conclusiones: Existe un papel mediador de los síntomas depresivos entre la exposición a violencia de pareja y el nivel de discapacidad en las mujeres del suroccidente colombiano. Así, tratar los síntomas de depresión es fundamental para mejorar la funcionalidad de las mujeres víctimas de la violencia de pareja. Los resultados de esta investigación contribuyen a definir estrategias para prevenir y abordar la violencia de pareja, los síntomas depresivos y la discapacidad en esta población.

© 2021 Asociación Colombiana de Psiquiatría. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Introduction

Violence is an important public health problem and one of the main causes of death worldwide. Each year, 1.7 million people lose their lives to violence and a considerable proportion suffer non-fatal injuries as a result of either self-inflicted, collective or interpersonal violence. Included in the latter is intimate partner violence (IPV), which is carried out by a member of a current or previous intimate relationship.¹ According to the World Health Organisation (WHO), 1 in every 3 women worldwide has experienced physical or sexual violence, which was predominantly conducted by their partner.² Out of the 50 000 registered cases of IPV in Colombia in 2018, 86% were against

women (43 000 cases), with a rate of 204/100 000 women. Cali, with 2100 cases, was the city with the second highest number of cases in Colombia.³

While the prevalence of IPV varies greatly between countries due to contextual factors, measuring IPV and comparing data across countries is also a challenge. Some of the differences in prevalence and the level of severity of IPV reported by women in different countries may be partly explained by the definitions of IPV used, the instruments used to measure it and the methodological designs of each study. For example, studies from around the world quote different prevalences of IPV: the United States of America, 64%;⁴ Egypt, North Africa, 29%;⁵ and Ghana, West Africa, 50%.⁶

IPV is a complex event that can vary depending on personal factors, such as age and character and on contextual factors which include situational and sociocultural factors.⁷ Contextual factors include family dynamics and economic dependence. Survivors of IPV also experience ill-health effects at an individual level and their functioning in society is impaired. Previous studies from different contexts have reported that IPV is associated with an increased risk of depressive symptoms, depression, PTSD and anxiety.⁸⁻¹²

IPV is a systematic behaviour that can be explained by the theory the cycle of abuse of IPV that consists of three interdependent phases: tension building, episode of aggression and the honeymoon. This violence has consequences on the survivors coping mechanisms associated with self-care, work, interpersonal relationships, and other areas.¹³

Mental health consequences of experiencing IPV can lead to deliberate self-harm and suicide.¹⁴⁻¹⁶ Unhealthy habits have also been observed such as alcohol abuse, smoking, the transmission of sexually transmitted diseases¹⁷ and physical injuries. A meta-analysis identified that the prevalence of depression in female survivors of IPV was approximately 48% and PTSD was between 24% and 84%.¹⁸

The 2015 Colombian National Survey of Mental Health reported that 9% of the 3463 women interviewed, aged 18 to 44 years old, had a traumatic experience due to domestic violence. Of these women, 57% stated the nature of it was psychological violence, 71% physical violence and 27% sexual violence. In women above 45 years old, 7% of 3023 subjects had a traumatic experience due to domestic violence and the figures were similar in relation to psychological (72%), physical (77%), and sexual violence (32%).¹⁹

IPV affects a women's quality of life as it is associated with different levels of disability; a concept defined by the WHO in its publication: The International Classification of Functioning, Disability and Health (ICF). They define disability as "an umbrella term for impairments, activity limitations and participation restrictions" and explain how "disability and functioning are viewed as outcomes of interactions between health conditions (diseases, disorders and injuries) and contextual factors". Contextual factors are divided into environmental (social attitudes, legal and social structures and climate, etc.) and personal (age, sex or ethnicity, etc.), which affect how disability is experienced.²⁰

Previous studies have identified that there is a relationship between depression and level of disability in female survivors of IPV.²¹⁻²³ In such cases, the disability is manifested at both societal and individual levels, taking into account the severity of violence. At the societal level it is associated with role loss and restrictions in participation, causing social isolation. At the individual level, it can be associated with experiencing limitations in conducting every day activities and both functional and physical impairments.^{24,25} While some studies have identified the close relationship between depression, IPV and disability, the mediating role of depression on the latter two is not clear.^{21,22} A mediating variable is one that transmits its effect on the relationship of an independent variable on a dependent variable.²⁶

Estimating the direct, indirect or total effect of an exposure on an outcome allows the identification of mediating effects between a group of variables.²⁶ Detecting mediation effects is

useful to identify casual pathways that generate a final outcome, in this case, disability. Targeted interventions can that be created to target the mediator, which in turn positively affects the outcome. The objective of the present study is to identify the mediating effect of depressive symptoms on the relationship between the severity of IPV experienced and the level of disability experienced in female survivors of IPV.

Methods

This was a cross-sectional study performed at the baseline of a randomised control trial "Evaluation of a cognitive-behavioural intervention for victims of domestic violence in Cali and Tuluá, Colombia", registered in ClinicalTrials.gov NCT03333798. This study recruited female adults (above 18 years old) during the period from November 14th 2017 to September 27th 2019. They were survivors of IPV performed by men who were identified by two public hospitals in the South West of Colombia. Women were excluded if they had severe pathologies or symptoms of mental illness, thyroid disease or if they had received pharmacological treatment for their mental health illnesses for less than 6 months.

Instruments

The level of severity of IPV in women was measured using the Severity of Violence Against Women Scale (SVAWS), that measured the violence they had experienced by their male partner within the last 12 months. The instrument uses 3 sub-scales: threats of violence (19 questions), physical violence (20 questions) and sexual violence (4 questions). SVAWS has shown a high reliability (Cronbach alpha, 0.89-0.96),²⁷ and it has been validated and applied previously in Colombia.²⁸ The questionnaire has the following responses to questions on the frequency of exposure to different violent behaviours: Never: 0 points; once: 1 point; sometimes: 2 points, and many times: 3 points. The total number of points give an overall score for each item on the SVAWS scale. There are total of 43 items with total possible scores ranging between 0 and 129.²⁸

For the analysis of the severity of violence, the scores of the three types of violence were combined and the answer to each item score was weighted, as previously done by Valdez-Santiago et al.,²⁹ where higher scores indicated higher severities of violence. Weighted scores ranged between 0 to 7, depending on the relative severity of the type of violence examined by each item. With this method, the total weighted score ranged from zero to 387 points, where a higher score indicated a higher severity of violence. Due to the wide range of values in the SVAWS weighted scale, their value was divided by 10 for the statistical analysis in the Structural Equation Model (SEM) described later.

The presence of depressive symptoms was evaluated using the instrument Hopkins Symptom Checklist (HSCL-25), which is a self-reporting scale, originally designed by Parloff et al.³⁰, later reduced to 25 questions.^{31,32} The HSCL-25 instrument has previously been validated and used in Colombia with a reported $\alpha=0.88$.³³ The HSCL-25 instrument asks about the frequency of depressive symptoms in the last four weeks and has the following responses: never: 1 point; once: 2 points;

sometimes: 3 points, and many times: 4 points. The score of the scale is calculated by taking the average score of all of the items and therefore has a range between 1 to 4 points, with 4 being the maximum score for depressive symptoms. Due to the narrow range of scores on this scale, the scores were multiplied by 10 for the analysis in the SEM. To measure the levels of disability in the study participants, the 12 item version of the Questionnaire for the Evaluation of Disability by the WHO (WHODAS 2.0) was used. This questionnaire enables an objective assessment of disability to be conducted that incorporates 6 life domains. These domains have been grouped into factors that limit activity and restrict participation and include: cognition, mobility, self-care, relationships, daily activities and participation.³⁴ In Colombia, the WHODAS 2.0 questionnaire has been applied in diverse populations,³⁵⁻⁴⁰ including survivors of IPV.⁴¹ This is a generic disability tool with response options relating to whether difficulty is experienced and if so, how much: no, zero points; mild, 1 point; moderate, 2 points; severe, 3 points, and extreme, 4 points. This questionnaire was validated in its original version and showed a temporal stability with an intraclass correlation coefficient of .69-.89 at the item level and .98 for the total score.^{42,43} The WHODAS 2.0 questionnaire scores are calculated through a complex method that establishes a scale from 0 (nil disability) to 100 points (complete disability).⁴⁴

Data collection

The study instruments were administered in face-to-face interviews conducted by 6 female health professionals: 4 psychologists, 1 registered nurse and 1 general physician who had all been trained in their use.

Statistical analysis

Bivariate analyses of the sociodemographic characteristics and depressive symptoms and disability levels respectively were performed using the Mann-Whitney U and Kruskal-Wallis tests. A path analysis using the SEM was used to evaluate the mediating role of depressive symptoms on the severity of violence and level of disability. Four observed exogenous variables were used in the model which had obtained $P < .20$ in the bivariate analysis: age, current relationship status (in a relationship/single), schooling (low: has not finished secondary school education, and high: has attained secondary school education or more) and preceding impairments (self-reported). Impairments are defined as "problems in body function or structure such as a significant deviation or loss".⁴⁵ The following endogenous variables were used in the model: severity of IPV and level of depressive symptoms (intermediate endogenous variables) and the level of disability (the main endogenous variable). The direct effect of severity of violence on disability was analysed as well as the indirect and mediating effect of depressive symptoms and the total effect of each explanatory variable.⁴⁶

The method of maximum likelihood was used for the estimation of the SEM, including missing values. The β coefficients of the SEM are presented in a non-standardised form to aid their interpretation (Table 2 and figure 1). The comparative fit index (CFI) $> .95$, the Tucker-Lewis Index (TLI) $> .95$, and

Table 1 – Sociodemographic characteristics of the study population.

<i>Age</i>	
Mean \pm SD	36 \pm 11
Median [ICR]	35 [26-43]
18-35	48 (52)
> 35-66	45 (48)
<i>Marital status</i>	
Single	41 (44)
Married	10 (11)
Separated	20 (21)
Cohabiting with partner	23 (24)
<i>Current relationship status</i>	
In a relationship	45 (48)
Single	49 (52)
<i>Education attainment</i>	
Low: nil schooling, primary school complete or incomplete, secondary school incomplete	40 (43)
High: secondary complete, technical certificate, university	54 (57)
<i>Employment status</i>	
Unemployed	37 (39)
Employed	57 (61)

the root mean square error (RMSE) $< .05$ were used to establish the model adjustment.^{46,47} All the analyses were performed in Stata[®] 14.2.

Results

Table 1 shows the sociodemographic characteristics of the study population. The median age was 35 [interquartile range, 26-43] years old, 41 women (44%) were married and 45 (48%) were in a relationship; 54 women (48%) had completed secondary education or had reached a technical or university level, and 57 (61%) were employed (Table 1).

The median violence weighted score reported by the participants was 104 [43-175] points, which was 27% of the highest obtainable score in the weighted SVWAS scale (points range, 0-387). The median depressive symptoms score was 2.8 [2-3] points, that corresponded to 60% of the highest obtainable score in the HSCL-25 instrument (points range, 1-4).

The median disability score of the participants was 28 [19-44], which can be interpreted as the percentage of disability experienced by responders to the WHODAS 2.0 questionnaire. There were no statistically significant differences between the median disabilities in each of the sociodemographic variables ($P > .05$). The median disability score for women aged 35 to 66 years old was 33 [19-48] in comparison with women aged 18 to 35, who had a median score of 27 [18-36]. In both cases these medians correspond to a moderate disability (25 to 49 points).

Figure 1 and Table 2 show the non-standardised coefficients of the causal pathways between severity of IPV, level of depressive symptoms and levels of disability. The SEM shows a good fit for the data according to the indicators (χ^2 , $P = .95$; CFI = 1.00; TLI = 1.23; RMSE < 0.01).

Figure 1 shows that low educational attainment was associated with an increase in the level of IPV experienced ($\beta = 4.28$; $P = .01$) as a direct effect. In contrast, having a partner and




Figure 1 – Casual path model and non-standardised coefficients. Boxes with dashed border contain the exogenous variables. Boxes with dotted border contain the intermediate endogenous variables, and the box with the continuous line contains the main endogenous variable. *Significant coefficients at $P < .05$.

a previous self-reported impairment were associated with experiencing lower levels of IPV ($\beta = -4.59$; $P = .005$ and $\beta = -6.50$; $P = .005$ respectively). In relation to depressive symptoms, they were found to be higher in women with a current partner ($\beta = 3.64$; $P = .003$) and with a self-reported previous impairment ($\beta = 4.57$; $P = .008$). All these results were statistically significant.

The results show that a mediating role exists between IPV severity and level of disability. The direct effect of IPV severity on level of disability was not statistically significant ($\beta = 0.09$; $P = .63$), however, the indirect effect of IPV severity on disability mediated by depressive symptoms was ($\beta = 0.39$; $P < .01$). The total effect of IPV severity on the level of disability was even greater ($\beta = 0.48$; $P = .01$). Additionally, the direct effect of depressive symptoms on the level of disability was also statistically significant ($\beta = 1.08$; $P < .01$).

With respect to the remaining total effects, depressive symptoms were positively affected by IPV severity ($\beta = 0.36$; $P < .01$) and low educational attainment ($\beta = 1.56$; $P = .02$). Finally, low educational attainment was marginally positively associated with level of disability ($\beta = 2.09$; $P = .07$).

Discussion

The objective of this study was to determine the mediating role of depressive symptoms on the relationship between the severity of IPV experienced and the level of disability in female survivors of IPV in Cali and Tuluá, Colombia.

The median weighted-score of IPV experienced by women in this study was 104 [42.5-175] points, which was 27% of the highest obtainable score of the weighted SVAWS scale. This median was similar to that reported by a previous study in Cali, where they found a mean violence of 41 ± 29 points in adult female survivors of IPV attending community centres. This mean value corresponded to 30% of the maximum score in the version of the SVAWS scale used (maximum, 138 points).²⁸ However, this study used the SVAWS instrument without considering weighted-scores for each of the items used in their research. This was advisable according to the method proposed by Valdez et al.,²⁹ which takes into account the severity of different expressions of violence.

The documented prevalence of depression among women who have experienced IPV varies, for example it has been reported as being 47% in North-American women¹⁸ and up to 75% in women in China.⁴⁸ The median score for the number of depressive symptoms in the HSCL-25 instrument reported by women in this study was 2.8 [2.4-3.4] out of a possible maximum of 4. The cut off point for positive cases of depression (1.75 points) was met or passed by an alarming 98% of women in Cali and 89% in Tuluá.

This study found that the direct effect of severity of IPV on the level of disability was not significant ($\beta = 0.09$; $P = .63$). However, it did find a statistically significant indirect affect and therefore mediating role of depressive symptoms on the relationship between the severity of IPV and the level of disability ($\beta = 0.39$; $P < .01$).

In view of these findings, this study suggests that after a woman is exposed to IPV, the presence of depressive symptoms is one of the main casual pathways associated with disability. In spite of the scarce amount of evidence that supports this type of analysis, there have been many reports of associations between the 3 different variables in this study. For instance, Bogantes-Rojas,²⁵ in their analysis on the cycle of abuse of domestic violence, describe depression and the presence of a physical or mental disability as direct consequences of IPV.

Both pre-existing physical and mental health illnesses, and those directly associated with IPV cause varying levels of disability. This study measured the level of disability using the WHODAS 2.0 questionnaire that evaluates disability taking a biopsychosocial approach with an emphasis on limitations on activity and restrictions on participation.⁴⁴ In the United States, a study investigated the impact of IPV on different aspects of "functioning", and found that nearly 45% of women with depressive symptoms had difficulties in their ability to work, attend school activities and participate in activities of daily living. They also reported difficulties in establishing effective social interactions in every day environments and in finding social support.²³

This data implies that depressive symptoms generate restrictions on the social participation of female survivors of IPV who are already facing difficult situations such as

Table 2 – Direct effects, indirect effects and total effects on the relationships between severity of IPV, depressive symptoms and levels of disability.

Parameters	Coefficient β	95%CI	P
Direct effects			
<i>Violence severity^a</i>			
Age ^b	-0.85	-2.33 to 0.63	.26
In a relationship (yes/no)	-4.59	-7.84 to -1.35	<.01
Low educational attainment (yes/no)	4.28	0.87 to 7.69	.01
Previous impairment (yes/no)	-6.50	-11.09 to -1.92	<.01
<i>Depressive symptoms^c</i>			
Violence severity ^a	0.36	0.22 to 0.50	<.01
In a relationship (yes/no)	3.64	1.25 to 6.03	<.01
Previous impairment (yes/no)	4.57	1.16 to 7.98	<.01
<i>Level of disability</i>			
Violence severity ^a	0.09	-0.28 to 0.46	.63
Depressive symptoms ^c	1.08	0.40 to 1.04	<.01
Age ^b	2.14	-0.39 to 4.69	.98
Previous impairment (yes/no)	4.99	1.75 to 11.97	.24
Indirect effects			
<i>Depressive symptoms^c</i>			
Age ^b	-0.30	-0.86 to 0.24	.27
In a relationship (yes/no)	-1.57	-1.20 to 0.37	.01
Low educational attainment (yes/no)	1.56	0.26 to 4.42	.02
Previous impairment (yes/no)	-2.37	-4.28 to -0.45	.15
<i>Level of disability</i>			
Violence severity ^a	0.39	0.24 to 0.55	<.01
Age ^b	-0.41	-1.20 to 0.37	.20
In a relationship (yes/no)	1.72	-2.02 to 0.46	.36
Low educational attainment (yes/no)	2.09	-0.21 to 4.41	.07
Previous impairment (yes/no)	1.80	-3.37 to 6.98	.49
Total effects			
<i>Violence severity^a</i>			
Age ^b	-0.84	-2.33 to 0.63	.26
In a relationship (yes/no)	-4.59	-7.84 to -1.35	<.01
Low educational attainment (yes/no)	4.28	0.87 to 7.69	.01
Previous impairment (yes/no)	-6.50	-11.09 to -1.92	<.01
<i>Depressive symptoms^c</i>			
Violence severity ^a	0.36	0.22 to 0.50	<.01
Age ^b	-0.46	-1.29 to 0.36	.27
In a relationship (yes/no)	2.95	-0.93 to 0.83	.13
Low educational attainment (yes/no)	1.56	0.17 to 2.94	.02
Previous impairment (yes/no)	2.20	-1.50 to 5.91	.24
<i>Level of disability</i>			
Violence severity ^a	0.48	0.08 to 0.89	.01
Depressive symptoms ^c	1.08	0.60 to 1.57	<.01
Age ^b	1.73	-0.91 to 4.38	.19
In a relationship (yes/no)	1.72	-2.02 to 5.46	.36
Low educational attainment (yes/no)	2.09	-0.21 to 4.42	.07
Previous impairment (yes/no)	6.80	-1.50 to 5.91	.14
95%CI: 95% confidence interval.			
^a IPV severity divided by 10.			
^b Age divided by 10.			
^c Depression scale multiplied by 10.			

social isolation and role loss.^{23,49} There are scarce studies from Colombia that evaluate the relationship between IPV, depressive symptoms and disability. Furthermore, theoretical models studying the relationships between health conditions and their context are not widely available.

This study found that depressive symptoms significantly, directly affected the level of disability, and mediated the effect of IPV severity on functional impairment levels. Similar results

were found in a longitudinal path analysis among women exposed to previous mass conflict traumas in post-conflict Timor-Leste, where both disability and IPV were associated with depressive symptoms. Authors have previously planted ideas that depression is related to one's self-perception of their physical and mental health and subjective experience of living with disability. This assumption substantiates the importance of performing multivariate analyses that contain the personal

and contextual factors (variables) in survivors of IPV in predictive models of disability.⁵⁰

Some findings in this current study were contradictory to those found in previous literature. Firstly, women in this study with an existing impairment were found to have experienced less severe levels of IPV. However, other studies have found that health conditions leading to disabilities increased the probability of having experienced physical, psychological and sexual IPV. This is thought to be due to these women being more vulnerable due to their condition and being economically dependent on their dominant partner.⁵¹

This study also found that women in this study who had a current partner had experienced lower levels of violence. As participants were asked about their experiences of violence within the last 12 months only, it may have not detected women who had experienced IPV by their ex-partners. A study from Mexico by Zaldivar-Cerón et al.⁵² found that separating from the aggressor does not imply an end to IPV by that same partner. On the contrary, it can lead to violent behaviour due to the threat that the perpetrator may lose control and power over their target. This behaviour can be explained by the cycle of abuse, where the honeymoon phase is not reached.

The sample size of this study is an important limitation that made it impossible to identify possible associations between exposures that were not considered in the SEM, for example alcohol and illicit drug abuse by the woman or her intimate partner. Given the complex phenomena of depression, IPV and disability, they have multiple explanatory variables. These are pertinent in understanding the pathways leading to them. Despite this limitation, the SEM was able to identify associations at the individual level, such as a previous impairment and education level as endogenous variables.

According to the authors' knowledge, the SEM analysis used in this study has proved to be innovative in Colombia where it has not previously been used in the context of depression, disability and IPV. Additionally, the SEM analysis enabled a simultaneous model to be run that included the study of relationships between multiple independent and dependent variables that are pertinent for the explanation of the complex phenomena of depression, IPV and disability.^{47,53}

Conclusions

The SEM showed that depressive symptoms have a mediating role on the relationship between the severity of exposure to IPV and the level of disability in women from 2 cities in the South West of Colombia. Alarming high levels of depression were found in this study which were also associated with experiencing IPV. Depression was also associated with having a self-reported impairment.

Education was an important additional factor found to be linked with both IPV and depression, which suggests that improved access to education is important in tackling these and ultimately disability.

Overall, these results indicate that attention to the amelioration of depressive symptoms are an essential step to improve the functionality of female survivors of IPV in middle income countries.

Financing

The authors thank the management and research teams of the CISALVA Institute of the Universidad del Valle (Cali, Colombia) and the Unidad Central del Valle del Cauca (-UCEVA- Tuluá, Colombia), who participated in the project "Evaluation of a Cognitive - Behavioural Intervention for victims of Intimate Partner Violence in Cali and Tuluá, Valle del Cauca", funded by MinCiencias (Colombia), Contract No. 756-2016. They also thank the women who participated in this study.

Conflict of interest

None.

REFERENCES

1. Krug EG, Dahlberg LL, Mercy JA, Zwi AB. Informe mundial sobre la violencia y la salud. SciELO Brasil; 2003.
2. World Health Organization. *Gender, equity and human rights* [Internet]. 2009. Available from: https://www.who.int/gender/violence/who_multicountry_study/summary_report/chapter2/es/.
3. Instituto Nacional de Medicina Legal y Ciencias Forenses. *Forensis*. 2018.
4. Thornton A, Young-DeMarco L. Four decades of trends in attitudes toward family issues in the United States: The 1960s through the 1990s. *J Marriage Fam*. 2001;63:1009-37.
5. Yaya S, Hudani A, Buh A, Bishwajit G. Prevalence and predictors of intimate partner violence among married women in Egypt. *J Interpers Violence*. 2019, 0886260519888196.
6. Alangea DO, Addo-Lartey AA, Sikweyiya Y, et al. Prevalence and risk factors of intimate partner violence among women in four districts of the central region of Ghana: Baseline findings from a cluster randomised controlled trial. *PLoS One*. 2018;13(7).
7. Heise LL. Violence against women: An integrated, ecological framework. *Violence Against Women*. 1998;4:262-90.
8. Lagdon S, Armour C, Stringer M. Adult experience of mental health outcomes as a result of intimate partner violence victimisation: a systematic review. *Eur J Psychotraumatol*. 2014;5:24794.
9. Wathen CN, MacGregor JCD, MacQuarrie BJ. Relationships among intimate partner violence, work, and health. *J Interpers Violence*. 2018;33:2268-70.
10. La Flair LN, Bradshaw CP, Campbell JC. Intimate partner violence/abuse and depressive symptoms among female health care workers: Longitudinal findings. *Women's Health Issues*. 2012;22:e53-9.
11. Devries KM, Mak JY, Bacchus LJ, et al. Intimate partner violence and incident depressive symptoms and suicide attempts: a systematic review of longitudinal studies. *PLoS Med*. 2013;10(5).
12. Castillo-Manzano RM, Arankowsky-Sandoval G. Violencia intrafamiliar como factor de riesgo para trastorno depresivo mayor en mujeres: Estudio de casos y controles. *Rev Biomedica*. 2008;19:128-35.
13. Llosa Martínez S, Canetti Wasser A. Depresión e ideación suicida en mujeres víctimas de violencia de pareja. *Psicol Conoc Soc*. 2019;9:138-60.
14. Ellsberg M, Emmelin M. Intimate partner violence and mental health. Taylor & Francis; 2014.

15. Rees S, Silove D, Chey T, et al. Lifetime prevalence of gender-based violence in women and the relationship with mental disorders and psychosocial function. *JAMA*. 2011;306:513-21.
16. Rivas-Rivero E, Bonilla-Algovia E. Salud mental y miedo a la separación en mujeres víctimas de violencia de pareja. *Rev Iberoam Psicol Salud*. 2020;11(1).
17. Hansen NB, Eriksen SB, Elklit A. Effects of an intervention program for female victims of intimate partner violence on psychological symptoms and perceived social support. *Eur J Psychotraumatol*. 2014;5:24797.
18. Golding JM. Intimate partner violence as a risk factor for mental disorders: A meta-analysis. *J Fam Violence*. 1999;14:99-132.
19. Gómez-Restrepo C. La Encuesta Nacional de Salud Mental-ENSM 2015. *Rev Colomb Psiquiatr*. 2016;4.
20. World Health Organization. International classification of functioning, disability and health: ICF. Geneva: World Health Organization; 2001.
21. Gibbs A, Carpenter B, Crankshaw T, et al. Prevalence and factors associated with recent intimate partner violence and relationships between disability and depression in post-partum women in one clinic in eThekweni Municipality. South Africa. *PLoS One*. 2017;12:e0181236.
22. Hanass-Hancock J, Carpenter B, Myezwa H. The missing link: exploring the intersection of gender, capabilities, and depressive symptoms in the context of chronic HIV. *Women Health*. 2019;59:1212-26.
23. Helfrich CA, Fujiura GT, Rutkowski-Kmitta V. Mental health disorders and functioning of women in domestic violence shelters. *J Interpers Violence*. 2008;23:437-53.
24. Silove D, Mohsin M, Klein L, et al. Longitudinal path analysis of depressive symptoms and functioning among women of child-rearing age in postconflict Timor-Leste. *BMJ Global Health*. 2020;5:e002039.
25. Bogantes Rojas J. Violencia doméstica. *Med Leg Costa Rica*. 2008;25:55-60.
26. VanderWeele T. Explanation in causal inference: methods for mediation and interaction. Oxford: Oxford University Press; 2015.
27. Marshall LL. Development of the severity of violence against women scales. *J Fam Violence*. 1992;7:103-21.
28. Burgos D, Canaval GE, Tobo N, Bernal de Pheils P, Humphreys J. Violencia de pareja en mujeres de la comunidad, tipos y severidad Cali, Colombia. *Rev Salud Pública*. 2012;14:377-89.
29. Valdez-Santiago R, Híjar-Medina MC, Salgado de Snyder VN, Rivera-Rivera L, Avila-Burgos L, Rojas R. Escala de violencia e índice de severidad: una propuesta metodológica para medir la violencia de pareja en mujeres mexicanas. *Salud Publica Mex*. 2006;48:s221-31.
30. Parloff MB, Kelman HC, Frank JD. Comfort, effectiveness, and self-awareness as criteria of improvement in psychotherapy. *Am J Psychiatry*. 1954;111:343-52.
31. Hesbacher PT, Rickels K, Morris RJ, Newman H, Rosenfeld H. Psychiatric illness in family practice. *J Clin Psychiatry*. 1980.
32. Winokur A, Winokur DF, Rickels K, Cox DS. Symptoms of emotional distress in a family planning service: stability over a four-week period. *Br J Psychiatry*. 1984;144:395-9.
33. Bonilla-Escobar FJ, Fandiño-Losada A, Martínez-Buitrago DM, et al. A randomized controlled trial of a transdiagnostic cognitive-behavioral intervention for Afro-descendants' survivors of systemic violence in Colombia. *PLoS One*. 2018;13:e0208483.
34. Vásquez-Barquero JL, Herrera S, Vásquez E, Gaité I. Cuestionario para la evaluación de discapacidad de la Organización Mundial de la Salud-WHO-DAS II (Versión española del World Health Organization Disability Assessment Schedule II). Madrid: Ministerio de Trabajo y Asuntos Sociales; 2006.
35. Henao C, Pérez J, Acosta Ojalora ML, Parra Sierra SJ, Quino Avila AC. Predicción del grado de discapacidad en adultos con lesión medular de Bogotá, usando el WHO-DAS II. *Rehabilitacion*. 2015;49:75-81.
36. Castro ES, Cargona LM, Bustos C, Pacheco Ro, Saldivia S. Modelo explicativo de discapacidad en población con trastornos mentales graves atendida en una institución especializada de la ciudad de Cali (Colombia) Explanatory Model of Disability in a Population with Severe Mental Disorders Attended in a Specializ. *Rev Ciencias Salud*. 2019;17:60-80.
37. Castro E, Castillo A. Factores relacionados con la carga de discapacidad en personas con intento de suicidio atendidos en la red pública de servicios de salud de la ciudad de Cali. Cali: Universidad del Valle. 2010.
38. Castaño A, Ceballos M, Vera C, Lugo LH. Funcionamiento y estado de salud en una población de amputados de miembro inferior en Medellín. Colombia. 2016;29:122-35.
39. Valencia F, Rivera D, Londoño J. Modelo predictivo del grado de discapacidad en adultos con lesión medular de Medellín. Antioquia. 2012.
40. Henao CP, Pérez JE. Modelo predictivo del grado de discapacidad en adultos con lesión medular: resultados desde el WHO-DAS II. *Rev Ciencias Salud*. 2011;9:159-72.
41. Ortiz M, Castillo A. Factores asociados a la ocurrencia de discapacidad física causada por lesiones intencionales correspondientes a violencia interpersonal atendidos en tres instituciones de salud de la ciudad de Cali. Cal: Universidad del Valle. 2011.
42. Abedzadeh M, Razi E, Sehat M. Psychometric properties of the world health organization disability assessment schedule II -12 Item (WHODAS II) in trauma patients. *Injury*. 2015.
43. Tarvonen-Schröder S, Tenovu O, Kaljonen A, Laimi K. Usability of World Health Organization disability assessment schedule in chronic traumatic brain injury. *J Rehabil Med*. 2018;50:514-8.
44. OMS, Servicio Nacional de Rehabilitación. Medición de la salud y la discapacidad manual para el cuestionario de evaluación de la discapacidad de la OMS WHODAS 2.0. Geneva: OMS; 2015.
45. World Health Organization. Towards a common language for functioning, disability and health ICF. *Int Classif*. 2002;1149:1-22.
46. Acock AC. Discovering structural equation modeling using Stata. Stata Press books. 2013.
47. Kaplan D. Structural equation modeling: Foundations and extensions. New York: Sage; 2008.
48. Wong JY-H, Tiwari A, Fong DY-T, Humphreys J, Bullock L. Depression among women experiencing intimate partner violence in a Chinese community. *Nurs Res*. 2011;60:58-65.
49. Ruiz-Pérez I, Rodríguez-Barranco M, Cervilla JA, Ricci-Cabello I. Intimate partner violence and mental disorders: co-occurrence and gender differences in a large cross-sectional population based study in Spain. *J Affect Disord*. 2018;229:69-78.
50. Dekel R, Shaked O, Ben-Porat A, Itzhaky H. The interrelations of physical and mental health: self-rated health, depression, and PTSD among female IPV survivors. *Violence Against Women*. 2020;26:379-94.
51. Alhusen JL, Bloom T, Anderson J, Hughes RB. Intimate partner violence, reproductive coercion, and unintended pregnancy in women with disabilities. *Disabil Health J*. 2019;100849.
52. Zaldívar-Cerón A, Gurrola-Peña GM, Balcázar-Nava P, Moysén-Chimal A, Esquivel-Santoveña EE. Las mujeres separadas de cara a la violencia de sus exparejas. Rumbo a su caracterización. *CienciaUAT*. 2015;10:83-92.
53. Kline RB. Principles and practice of structural equation modeling. London: Guilford; 2015.