

Chemical composition and biological evaluation of tea tree (*Melaleuca alternifolia* L.) leaves essential oils

Composición química y evaluación biológica de los aceites esenciales de las hojas del árbol del té (*Melaleuca alternifolia* L.)

<https://doi.org/10.15446/rfnam.v77n2.109468>

Pham Thi Quyen¹ and Le Pham Tan Quoc^{1*}

ABSTRACT

Keywords:

Antioxidant capacity
Antibacterial activity
Essential oils
Herbs
GC-MS



Melaleuca alternifolia L. essential oils (MaEO), extracted through steam, distillation of its leaves, offer a multitude of benefits. The aim of the study was to determine the physicochemical and biological properties, such as relative density, absolute density, acid value, saponification value, ester value, freezing point, fragrance retention, antioxidant and antibacterial activity of MaEO. Gas chromatography-mass spectrometry (GC-MS) analysis was used to analyze the chemical composition of essential oils (EO), and the obtained results displayed those 45 compounds were identified and quantified; among them, the main component was Terpinen-4-ol (44.55%). In addition, the antioxidant capacity (AC) was evaluated by the 2,2-diphenyl-1-picrylhydrazyl (DPPH) assay with the half maximal inhibitory concentration (IC₅₀) of 360 mg mL⁻¹. Moreover, MaEO also had excellent antibacterial activity (AA) against certain types of bacteria, using the paper disc diffusion method for antibiotic susceptibility testing. The diameter of the inhibitory zones were 12.33, 14, 15.67, and 24.33 mm for *Bacillus cereus* (ATCC 11778), *Staphylococcus aureus* (ATCC 25923), *Salmonella enterica* (ATCC 13076), and *Escherichia coli* (ATCC 25922), respectively.

RESUMEN

Palabras clave:

Capacidad antioxidante
Actividad antibacteriana
Aceites esenciales
Hierbas
GC-MS

Los aceites esenciales de *Melaleuca alternifolia* L. (MaEO), extraídos mediante destilación al vapor de sus hojas, ofrecen múltiples beneficios. El objetivo del estudio fue determinar las propiedades fisicoquímicas y biológicas, como la densidad relativa, la densidad absoluta, el índice de acidez, el índice de saponificación, el índice de éster, el punto de congelación, la retención de fragancia, la actividad antioxidante y antibacteriana del MaEO. Se utilizó análisis de cromatografía de gases-espectrometría de masas (GC-MS) para analizar la composición química de los aceites esenciales (AE), y los resultados obtenidos mostraron que se identificaron y cuantificaron 45 compuestos; entre ellos, el componente principal fue terpinen-4-ol (44,55%). Además, la capacidad antioxidante (CA) se evaluó mediante el ensayo de 2,2-difenil-1-picrilhidrazilo (DPPH) con la concentración inhibitoria media máxima (IC₅₀) de 360 mg mL⁻¹. Asimismo, MaEO también tuvo una excelente actividad antibacteriana (AA) contra ciertos tipos de bacterias, utilizando el método de difusión en disco de papel para pruebas de susceptibilidad a los antibióticos. El diámetro de las zonas inhibitorias fueron 12,33, 14, 15,67 y 24,33 mm para *Bacillus cereus* (ATCC 11778), *Staphylococcus aureus* (ATCC 25923), *Salmonella enterica* (ATCC 13076) y *Escherichia coli* (ATCC 25922), respectivamente.

¹Institute of Biotechnology and Food Technology, Industrial University of Ho Chi Minh City, Ho Chi Minh City, Vietnam. phamthiquyen@iuh.edu.vn , lephamtanquoc@iuh.edu.vn 

*Corresponding author

M*elaleuca alternifolia* L., commonly known as the tea tree, belongs to the family Myrtaceae, native to Australia (Rodney et al. 2015). Currently, it is widely distributed globally, mainly in Australia, tropical America, South Asia, and Indonesia (Yasin et al. 2021). Tea tree is a white-barked shrub up to 7 m tall, with narrow leaves scattered into cymbals, 10–35 mm long and 1 mm wide, and the leaves are especially rich in essential oils. Tea tree flowers are white, each flower solitary in a bract, with petals 2–3 mm long. The fruit is a woody, cup-shaped capsule 2–3 mm in diameter and has many seeds (Rodney et al. 2015).

The leaves of this plant can help fight wound infections and are used in traditional medicine (Rodney et al. 2015; Yasin et al. 2021). *Melaleuca alternifolia* leaves essential oil (MaEO) has many applications in various fields, such as agriculture, pharmaceuticals, cosmetics, veterinary products, and food. Currently, tea tree growth is quite popular in Vietnam. Essential oils (EO) comprise terpene hydrocarbons, mainly monoterpenes, sesquiterpenes, and their related alcohols (Carson et al. 2006). According to ISO 4730 (2017), Terpinen-4-ol is the main ingredient in tea tree oil, with concentrations ranging from 35 to 48%. The ingredients' content significantly differs depending on geographical location, extraction method, harvesting season, and storage conditions (Borotová et al. 2022). MaEO has high biological activity, including antibacterial, antifungal, and antiviral activities, which has been studied on many bacteria and fungi (Carson et al. 2006). The application of MaEO in food preservation has been studied quite a lot in recent years. Typically, chitosan films obtained in malic acid with MaEO have the highest antioxidant activity, color, and insignificant transparency change (Cázon et al. 2021). In addition, MaEO is used to inhibit molds isolated from meat products (Sevik et al. 2021).

It knows that the same plant material distributed in different regions will have different characteristics. Until now, many studies have been related to the tea tree and its essential oil. However, this is the first time *M. alternifolia* has been grown in Lam Dong province (Vietnam) and the MaEO may differ from that in other regions regarding volatile chemical composition and biological activities. Therefore, this study aimed to evaluate the physicochemical properties, chemical profile, and biological activities of MaEO. The results of this study could contribute significantly to understanding this

essential oil in various regions. With its distinctive chemical composition and biological properties, this material holds potential applications in food technology, medicine, and cosmetics.

MATERIALS AND METHODS

Materials

The leaves of *M. alternifolia* were harvested after about 9 months of age in Lam Dong province, Vietnam (Coordinates: 11°40'1.20"N, 107°19'58.80"E). The plant specimen (coded MA151022VST) has been archived at the Plant Biotechnology Laboratory of the Institute of Biotechnology and Food Technology, Ho Chi Minh City University of Industry. On average, the per-batch yield was approximately 50 kg leaves/batch and MaEO was extracted using steam distillation for 3 h at 100 °C, with the yield obtained at about 1.5% (v/w), and the EO was stored in a dark sealed bottle at 4 °C before analysis.

Bacterial strains used in this study include *Staphylococcus aureus* (ATCC 25923), *Bacillus cereus* (ATCC 11778), *Escherichia coli* (ATCC 25922), and *Salmonella enterica* (ATCC 10376).

Determination of physicochemical properties of MaEO

The freezing point (FP), relative density (RD), absolute density (AD), acid value (AV), esters value (EV), and saponification value (SV) were evaluated according to International Organization for Standardization including ISO 1041 (1973), 279 (1998), 1242 (2023), and 7660 (1983), respectively.

Determination of fragrance retention (FR) of MaEO

According to the procedure described by Mahajan (2022), fragrance retention (FR) is determined by the concentration of the flavoring ingredient and its retention time, with some minor corrections. EO was mixed into concentrations (5, 10, 15, 20, and 25%, v/v) in 96% ethanol. Next, three drops of EO were placed on the odor test paper, and a few seconds were allotted for the EO to penetrate the paper. Finally, the time until the smell of EOs disappeared was calculated and the results recorded.

Gas chromatography-mass spectrometry (GC-MS)

The chemical volatile composition of EO was analyzed using the GC-MS method. 1 µL of EO was injected into a gas chromatograph (Shimadzu Nexis GC-2030, Japan)

with a versatile capillary column (Rtx-5sil-MS, 30 m×0.25 mm×0.25 µm, Restek Technologies, USA) equipped with a quadrupole mass analyzer (Shimadzu GC-MS-QP2020 NX, Japan). Helium was used as a carrier gas at a constant flow rate of 3 mL min⁻¹, and a split ratio of 10:1. The injection temperature was 250 °C and the temperature program was set as follows: initial temperature of 50 °C, held for 2 min, increased until 250 °C at a rate of 10 °C min⁻¹, and held for 5 min; finally, increased to 280 °C at a rate of 10 °C min⁻¹, and held for 3 min. Mass spectra were recorded at the ionization energy of 70 eV in EI mode (Hao and Quoc 2024).

Determination of antioxidant capacity (AC) of MaEO

The AC of MaEO was determined using DPPH assay

$$\% \text{ DPPH}_{\text{RSC}} = \frac{\text{Absorbance of control} - \text{Absorbance of sample}}{\text{Absorbance of control}} \times 100 \quad (1)$$

Determination of the antibacterial activity (AA) of MaEO

The AA was determined by susceptibility testing by the agar plate diffusion method described by Quyen and Quoc (2023) with minor adjustments. First, a sterile spreader spread 100 µL of bacterial suspension (0.5 McFarland standard, approximately 1.5×10⁸ CFU mL⁻¹) onto Muller-Hinton agar medium (MHA). Then, paper plates with a diameter of 6 mm were impregnated with essential oils (5 µL) put on MHA media, while ampicillin (10 µg disc⁻¹) and 5% dimethyl sulfoxide (DMSO) solution (5 µL disc⁻¹) were used as a positive and negative control. Petri dishes were incubated for 24 h at 37 °C, and the diameter of the inhibition zone was measured in mm.

Statistical data analysis

All the experimental results were analyzed using Statgraphics Centurion software (Version 15.1.02). Every assay was done in triplicates. Analysis of variance (ANOVA) with Fisher's least significant difference procedure was used to determine the significant differences ($P < 0.05$) between means.

RESULTS AND DISCUSSION

Determination of physicochemical properties of *Melaleuca alternifolia* leaf essential oils

MaEO is a light-yellow liquid with a characteristic odor and a bitter taste. The physicochemical properties of EO are shown in Table 1. The pH value of MaEO is about 4.94

according to the procedure described by Rahman et al. (2015) with some minor corrections. EOs were diluted into concentrations (100, 200, 300, 400, 500, and 600 mg mL⁻¹) in 96% ethanol. 0.3 mL of the obtained EO and 3.7 mL DPPH 0.1 M in 96% ethanol solution were mixed. The mixture was kept in dark conditions for 30 min at room temperature. DPPH radical scavenging capacity (DPPH_{RSC}) of MaEO was expressed by the degree of color reduction of the DPPH solution, as determined by measuring the absorbance at a wavelength of 517 nm, and antioxidant capacity was calculated following equation 1, while vitamin C was used as a control. Percent inhibition plotted against EO concentration to estimate concentration gives 50% inhibition (IC₅₀).

and this result is higher than that in some EOs from other materials, such as *Melaleuca cajuputi* leaves (pH=4.46) (Quoc 2021) and *Ceratonia siliqua* pulp (pH=4.3) (Ouis and Hariri 2018). The chemical component of EO affects pH strongly, leading to a significant difference in pH in materials.

The FP of EO remains undetermined at -40 °C; the oil still exists in liquid form, with no freezing phenomenon. The results show that the FP of MaEO must be below -40 °C; compared to other materials, FPs of the EO of *M. cajuputi* leaves and *Ocimum gratissimum* were -45 and -38 °C (Quoc 2021; Hao and Quoc 2024). This indicated that FP depends on the chemical composition of the EO significantly. In addition, the RD and AD of the oil were also analyzed; both values are lower than one, 0.9064 and 0.9048 g mL⁻¹, respectively. These findings agree with the ISO 4730 (2017), and they did not differ significantly from those of *M. cajuputi* (RD: 0.9102 and AD: 0.9086 g mL⁻¹) (Quoc 2021) or *M. arvensis* (RD: 0.8987 and AD: 0.8959 g mL⁻¹) leaves' EO (Quoc 2022). The obtained values also revealed that the MaEO belongs to the EO group lighter than water.

The AV, SV, and EV of MaEO were determined to be 2.10, 7.34, and 5.24 mg KOH g⁻¹ EO, respectively. In general, the AV is agreement with that of previous studies for other materials, such as *C. siliqua* pulp and seeds (AV:

3.82 and 2.2 mg g⁻¹ EO) (Ouis and Hariri 2018), basil EO (AV: 3.95 mg KOH g⁻¹ EO), and lemongrass (AV: 4.09 mg KOH g⁻¹ EO) (Mustapha 2018), while the obtained SV is significantly lower than that of *M. cajuputi* leaves EO (SV: 28.05 mg KOH g⁻¹ EO) (Quoc 2021), basil EO (SV: 198 mg

KOH g⁻¹ EO), and lemongrass (SV: 143 mg KOH g⁻¹ EO) (Mustapha 2018). The AV, SV, and EV are indicators used to assess the quality of the EO and they are influenced by distillation technique, cultivar, climatic conditions, harvest period, and chemical composition of the initial materials.

Table 1. Physicochemical properties of *Melaleuca alternifolia* leave essential oils.

No.	Physicochemical properties	Value
1	pH	4.94±0.09
2	Freezing point (FP, °C)	<-40
3	Relative density (RD)	0.9064±0.0015
4	Absolute density (AD, g mL ⁻¹)	0.9048±0.0015
5	Acid value (AV, mg KOH g ⁻¹ EO)	2.10±0.1354
6	Saponification value (SV, mg KOH g ⁻¹ EO)	7.34±0.8609
7	Ester value (EV, mg KOH g ⁻¹ EO)	5.24±0.8068
8	Fragrance retention (FR, h):	
	5% EO	1.25
	10% EO	2.17
	15% EO	5.42
	20% EO	6.17
	25% EO	10.42
	100% EO	25.25

The staying power is also the critical index to evaluate the quality of EO. In this study, FR of MaEO can reach a range of 6–10 h at 20–25%, and this shows that FR is quite long, similar to perfume at a fragrance concentration of 20–30%, and it can last for more than 6–8 h (Mahajan 2022), while pure MaEO can last more than 24 h. This finding proves that MaEO has potential applications in cosmetics and the food industry.

Chemical composition of *Melaleuca alternifolia* leaves essential oils

The volatile compounds of MaEO were analyzed by GC-MS. The results are displayed in Table 2. A total of 45 compounds were identified and quantified in the EO isolated from the *M. alternifolia* leaves; they account for 99.35% of the oil and are tested by retention times ranging from 7 to 22 min. Compounds comprising the highest content in MaEO, include Terpinen-4-ol (44.55%), γ -Terpinene (19.42%), p-Cymene (8.75%), α -Terpinene (6.73%), and Terpineol (3.33%). These components play an important role in the quality of MaEO and there is variation in the

chemical composition of EO from tea tree leaves collected in different regions. In previous studies, authors also used the distillation method to extract EOs from the initial material. For example, MaEO isolated from Tien Giang province (Vietnam) possess 19 volatile components: Terpinen-4-ol (36%), γ -Terpinene (17.8%), 1,8-Cineole (10%), etc. (Hòa et al. 2016), while MaEO from Slovakia has 47 identified components, including Terpinen-4-ol (40.3%), γ -Terpinene (11.7%), 1,8-Cineole (7%), p-Cymene (6.2%), etc. (Borotová et al. 2022), and while MaEO collected in Thailand has 24 components, including Terpinen-4-ol (30.42–34.76%), γ -Terpinene (25.08–26.23%), α -Terpinene (12.31–12.43%) and 1,8-Cineole (5.99–9.08%), etc. (Sukatta et al. 2011). In this study, the main component of MaEO is Terpinen-4-ol and its content is higher than that in MaEO from other places. The MaEO from the Lam Dong province complied with the ISO 4730 (2017) standard analysis with few variations in the contents of major constituents. This difference in the chemical composition of MaEO can be explained due to different extraction methods, climatic conditions, harvesting periods, genes, etc.

Table 2. Chemical composition of *M. alternifolia* leaves essential oils.

No.	Compounds	Molecular formula	RT. (min)	Content (%)
1	Bicyclo[3.1.0]hex-2-ene,2-methyl-5-(1-methylethyl)-	C ₁₀ H ₁₆	6.998	0.65
2	α-Pinene	C ₁₀ H ₁₆	7.146	2.35
3	Camphene	C ₁₀ H ₁₆	7.484	0.02
4	Ethanone	C ₈ H ₁₄ O	7.828	0.01
5	Bicyclo[3.1.0]hexane,4-methylene-1-(1-methylethyl)-	C ₁₀ H ₁₆	7.956	0.14
6	Bicyclo[3.1.1]heptane,6,6-dimethyl-2-methylene-, (1S)-	C ₁₀ H ₁₆	8.051	0.51
7	3-Methyl-3-cyclohexen-1-one	C ₇ H ₁₀ O	8.196	0.05
8	β-Myrcene	C ₁₀ H ₁₆	8.283	0.52
9	Dodecane	C ₁₂ H ₂₆	8.502	0.05
10	α-Phellandrene	C ₁₀ H ₁₆	8.612	0.29
11	α-Terpinene	C₁₀H₁₆	8.821	6.73
12	p-Cymene	C₁₀H₁₄	8.960	8.75
13	Bornylene	C ₁₀ H ₁₆	9.059	1.02
14	β-Phellandrene	C ₁₀ H ₁₆	9.085	0.41
15	1,8-Cineole	C ₁₀ H ₁₈ O	9.115	0.91
16	γ-Terpinene	C₁₀H₁₆	9.599	19.42
17	Cyclohexene	C ₁₀ H ₁₆	10.101	3.2
18	Benzene, (2-methyl-1-propenyl)-	C ₁₀ H ₁₂	10.169	0.15
19	Linalool	C ₁₀ H ₁₈ O	10.320	0.05
20	p-Mentha-1,5,8-triene	C ₁₀ H ₁₄	10.577	0.04
21	2-Cyclohexen-1-ol,1-methyl-4-(1-methylethyl)-, cis-	C ₁₀ H ₁₈ O	10.780	0.18
22	β-Ocimene	C ₁₀ H ₁₆	10.971	0.13
23	2-Cyclohexen-1-ol,1-methyl-4-(1-methylethyl)-, trans-	C ₁₀ H ₁₈ O	11.094	0.14
24	Benzene, (methyl(1-methylethyl)-	C ₁₀ H ₁₆ O	11.623	0.07
25	Terpinen-4-ol	C₁₀H₁₈O	11.745	44.55
26	p-Cymen-8-ol	C ₁₀ H ₁₄ O	11.815	0.04
27	Terpineol	C₁₀H₁₈O	11.968	3.33
28	p-Menth-2-en-1,4-diol	C ₁₀ H ₁₈ O ₂	13.224	0.3
29	cis,cis-Photocitral A	C ₁₀ H ₁₆ O	14.038	0.35
30	2-Cyclohexyl-hex-5-en-2-ol	C ₁₂ H ₂₂ O	14.845	0.03
31	α-Copaene	C ₁₅ H ₂₄	15.254	0.15
32	α-Gurjunene	C ₁₅ H ₂₄	16.037	0.31
33	Caryophyllene	C ₁₅ H ₂₄	16.363	0.25
34	Aromandendrene	C ₁₅ H ₂₄	16.886	0.97
35	Alloaromadendrene	C ₁₅ H ₂₄	17.529	0.31
36	δ-Cadinene	C ₁₅ H ₂₄	17.869	1.33
37	β-Selinene	C ₁₅ H ₂₄	18.435	0.1
38	1H-Xycloprop[e]azulen, 1a,2,3,5,6,7,7a,7b-octahydro-1,1,4,7-tetrametyl-	C ₁₅ H ₂₄	18.503	0.89
39	Naphtalen, 1,2,3,4-tetrahydro-1,6-dimetyl-4-(1-metyletyl)-, (1S-cis)-	C ₁₅ H ₂₂	19.305	0.21
40	Isolodene	C ₁₅ H ₂₄	19.365	0.23
41	Naphtalen,1,2,3,4,4a,7-hexahydro-1,6-dimetyl-4-(1-metyletyl)-	C ₁₅ H ₂₄	19.607	0.18
42	(-)-Globulol	C ₁₅ H ₂₆ O	20.874	0.21
43	Guaiol	C ₁₅ H ₂₆ O	21.103	0.23
44	Di-epi-1,10-cubenol	C ₁₅ H ₂₆ O	21.725	0.18
45	Epicubenol	C ₁₅ H ₂₆ O	22.018	0.06

Antioxidant capacity (AC) of *Melaleuca alternifolia* leaves essential oils

Figures 1 and 2 show the efficiency of the antioxidant process of EO and control (vitamin C). An increase in MaEO concentration leads to an increase in AC. Based on the obtained curve, the IC_{50} of MaEO was determined at 360 mg mL^{-1} (Figure 1), while the IC_{50} of vitamin C was only about $26 \mu\text{g mL}^{-1}$ (Figure 2). The results show that the IC_{50} of MaEO is much higher than that of vitamin C, showing that the AC of EO is very weak. These results

are consistent with those of MaEO from Thailand (IC_{50} : $29.34\text{--}38.68 \text{ mg mL}^{-1}$) (Sukatta et al. 2011) (Their AC is extremely low compared to the control). Besides, compared to some EOs from other materials, such as *C. siliqua* pulp and seeds (IC_{50} : 7.8 and $31.25 \mu\text{g mL}^{-1}$) (Ouis and Hariri 2018), *M. arvensis* (IC_{50} : 330 mg mL^{-1}) (Quoc 2022), the AC of MaEO is also weaker. This is a disadvantage when applying this EO in the cosmetic and food industry. Differences in AC of various EOs may be due to differences in their chemical composition.




Figure 1. Antioxidant capacity of the *Melaleuca alternifolia* leaves EO.




Figure 2. Antioxidant capacity of vitamin C.

Antibacterial activity (AA) of *Melaleuca alternifolia* leaves essential oils

Table 3 shows that MaEO exhibited to AA against four bacterial strains, and ampicillin was used as an inhibitory control against the bacteria used in the study. AA of

ampicillin are arranged in order of susceptibility: *E. coli* > *S. enterica* > *S. aureus* / *B. cereus*. Similarly, AA of MaEO is arranged in order of susceptibility: *E. coli* > *S. enterica* / *S. aureus* / *B. cereus*. The AA of MaEO was 1.5 times stronger than that of ampicillin for *S. aureus* and *B. cereus*, while

the AA of MaEO was much lower than that of ampicillin for *S. enterica* and *E. coli*. These obtained results are also different from those of MaEO collected from Thailand (Sukatta et al. 2011) and other materials (*C. siliqua* pulp and seeds, *M. cajuputi* leaves, and *O. gratissimum* EO) (Ouis and Hariri 2018; Quoc 2021; Hao and Quoc 2024). The AA of MaEO was observed in the presence of a high content

of Terpinen-4-ol; according to Halcón and Milkus (2004), Terpinen-4-ol strongly influences the bacterial cell wall and compromises the cytoplasmic membrane of bacteria (*S. aureus*), giving it a bacteriostatic and bactericidal effect. In addition, combining other bioactive ingredients, such as γ -Terpinene, 1,8-Cineole, α -Terpineol, etc., can also increase the ability to inhibit bacteria.

Table 3. Antibacterial zones of *M. alternifolia* leaves essential oils.

No.	Microorganisms	Diameter of the inhibitory zones of ampicillin (mm)	Diameter of the inhibitory zones of EO (mm)
1	<i>S. aureus</i>	9.33 ^{Aa} ±1.53	14 ^{Ba} ±1
2	<i>B. cereus</i>	8.33 ^{Aa} ±1.53	12.33 ^{Ba} ±2.31
3	<i>S. enterica</i>	22.67 ^{Bb} ±1.16	15.67 ^{Aa} ±1.16
4	<i>E. coli</i>	29.33 ^{Bc} ±4.04	24.33 ^{Ab} ±3.51

Within a row (A–B) or a column (a–c), different letters denote significant differences ($P < 0.05$) between samples or microorganisms, respectively.

CONCLUSION

The essential oil distilled from *M. alternifolia* leaves in Lam Dong province, Vietnam, exhibits unique physicochemical properties, notably exceptional fragrance retention lasting approximately 25 h for pure EO. It has weak AC compared to the control. However, this material also possesses high AA and strongly inhibits four pathogenic bacterial strains in food, including *E. coli*, *S. enterica*, *S. aureus*, and *B. cereus*. By the GC-MS method, 45 major volatile components were determined in tea tree leaf EO. This is a natural, precious, and rich source of phytochemicals, especially Terpinen-4-ol (44.55%), γ -Terpinene (19.42%), etc. Therefore, the oil from *M. alternifolia* leaves is ideal for use in the pharmaceutical, food, and cosmetic industries.

ACKNOWLEDGMENTS

This study was conducted at the Institute of Biotechnology and Food Technology, Industrial University of Ho Chi Minh City (Vietnam). The author would like to thank Le Van Hai and Tran Ngoc Kim Hien for their helpful advice on various technical issues considered in this article.

REFERENCES

Borotová P, Galovičová L, Vukovic NL et al (2022) Chemical and biological characterization of *Melaleuca alternifolia* essential oil. *Plants* 11(4): 558. <https://doi.org/10.3390/plants11040558>

Cázon P, Antoniewska A, Rutkowska J and Vázquez M (2021) Evaluation of easy-removing antioxidant films of chitosan with *Melaleuca alternifolia* essential oil. *International Journal of*

Biological Macromolecules 186: 365-376. <https://doi.org/10.1016/j.ijbiomac.2021.07.035>

Carson CF, Hammer KA and Riley TV (2006) *Melaleuca alternifolia* (tea tree) oil: a review of antimicrobial and other medicinal properties. *Clinical Microbiology Reviews* 19(1): 50-62. <https://doi.org/10.1128/cmr.19.1.50-62.2006>

Halcón L and Milkus K (2004) *Staphylococcus aureus* and wounds: A review of tea tree oil as a promising antimicrobial. *American Journal of Infection Control* 32(7): 402-408. <https://doi.org/10.1016/j.ajic.2003.12.008>

Hao PM and Quoc LPT (2024) Chemical profile and antimicrobial activity of *Ocimum gratissimum* L. essential oil from Dak Lak province, Vietnam. *Journal of Plant Biotechnology* 51: 50-54. <https://doi.org/10.5010/JPB.2024.51.005.050>

Hòa DM, Duy VH and Chi NTD (2016) Research on extraction, chemical compositions, and application of tea tree (*Melaleuca alternifolia*) essential oil in the production of mouthwash. *Can Tho University Journal of Science* 45: 90-96. <https://doi.org/10.22144/ctu.jvn.2016.515>

ISO – International Organisation for Standardisation (2017) Essential oil of *Melaleuca*, terpinen-4-ol type (Tea Tree oil). ISO 4730. Switzerland.

ISO – International Organization for Standardization (1973) Essential oils - Determination of freezing point. ISO 1041. Switzerland.

ISO – International Organization for Standardization (1983) Essential oils - Determination of ester value of oils containing difficult-to-saponify esters. ISO 7660. Switzerland.

ISO – International Organization for Standardization (1998) Essential oils - Determination of relative density at 20°C. ISO 279. Switzerland.

ISO – International Organization for Standardization (2023) Essential oils - Determination of acid value. ISO 1242. Switzerland.

Mahajan VK (2022) Perfumes and associated allergens: A brief review. *CosmoDerma* 2(21): 1-12. https://doi.org/10.25259/CSDM_9_2022

Mustapha A (2018) Comparative analysis on the extraction of essential oil from lemongrass and basil leaves. *International Journal of*

of Innovative Science, Engineering & Technology 5(11): 114-118.

Ouis N and Hariri A (2018) Antioxidant and antibacterial activities of the essential oils of *Ceratonia siliqua*. Banat's Journal of Biotechnology IX(17): 13-23.

Quoc LPT (2021) Physicochemical properties, chemical components, and antibacterial activity of *Melaleuca cajuputi* Powell essential oil leaves from Quang Tri province, Vietnam. Bulletin of the Chemical Society of Ethiopia 35(3): 677-683. <https://doi.org/10.4314/bcse.v35i3.18>

Quoc LPT (2022) Physicochemical properties, chemical components, and antibacterial activity of the essential oil from *Mentha arvensis* L. leaves. Uchenye Zapiski Kazanskogo Universiteta. Seriya Estestvennye Nauki 164(1): 36-45.

Quyen PT and Quoc LPT (2023) Chemical profile and biological properties of the essential oil of Rosemary leaves (*Rosmarinus officinalis* L.). Mongolian Journal of Chemistry 24(50): 40-45. <https://doi.org/10.5564/mjc.v24i50.2853>

Rahman MM, Islam MB, Biswas M and Alam AHMK (2015) In

vitro antioxidant and free radical scavenging activity of different parts of *Tabebuia pallida* growing in Bangladesh. BMC Research Notes 8: 621. <https://doi.org/10.1186/s13104-015-1618-6>

Rodney J, Sahari J and Shah MKM (2015) Review: Tea tree (*Melaleuca alternifolia*) as a new material for biocomposites. Journal of Applied Science and Agriculture 10(3): 21-39.

Sevik R, Akarca G, Kilinc M and Ascioğlu C (2021) Chemical composition of tea tree (*Melaleuca alternifolia*) (maiden & betche) cheel essential oil and its antifungal effect on foodborne molds isolated from meat products. Journal of Essential Oil Bearing Plants 24(3): 561-570. <https://doi.org/10.1080/0972060X.2021.1942232>

Sukatta U, Rughtaworn P, Tuntawiroon O and Meaktrong W (2011) Physico-chemical properties, chemical composition and *in vitro*. Kasetsart Journal - Natural Science 45: 473-480.

Yasin M, Younis A, Javed T et al (2021) River tea tree oil: composition, antimicrobial and antioxidant activities, and potential applications in agriculture. Plants 10(10): 2105. <https://doi.org/10.3390/plants10102105>